



# Clinical trials appendix

## Q1 2022



# Movement since Q4 2021 update

New to Phase I	New to Phase II	New to Pivotal trial	New to registration
<p><b><u>NME</u></b> <b>NI006<sup>#</sup></b> TTR deplete transthyretin amyloid cardiomyopathy</p>	<p><b><u>NME</u></b> <b>AZD4573</b> CDK9 inhibitor haematological malignancies</p>	<p><b><u>NME</u></b> <b>tozorakimab</b> IL-33 mAb COPD</p> <p><b><u>Additional indication</u></b> <b>Imfinzi + treme +TACE +/- lenvatinib EMERALD-3</b> PD-L1 + CTLA4 + VEGF +/- chemo-embolisation locoregional hepatocellular carcinoma</p> <p><b><u>Life-cycle management</u></b> <b>Fasenra HUDSON</b> IL-5R mAb eosinophilic gastritis and eosinophilic gastroenteritis</p> <p><b>Imfinzi + domvanalimab (AB154) + CTx PACIFIC-8<sup>#</sup></b> PD-L1 mAb + TIGIT + CTx Unresectable Stage III NSCLC</p> <p><b>Imfinzi +/- oleclumab +/- mona PACIFIC-9<sup>#</sup></b> PD-L1 + NKG2a or PD-L1 + CD73 Unresectable stage III NSCLC</p>	<p><b><u>NME</u></b> <b>nirsevimab<sup>#</sup></b> RSV mAb-YTE passive RSV immunisation</p> <p><b>Imfinzi + tremelimumab<sup>#</sup></b> PD-L1+CTLA-4 1L HCC HIMALAYA</p> <p><b><u>Life-cycle management</u></b> <b>Enhertu DESTINY-Lung01<sup>¶</sup></b> HER2 targeting antibody drug conjugate HER2-over-expressing or -mutated, unresectable and/or metastatic non-small cell lung cancer</p> <p><b>Imfinzi + CTx <sup>#</sup></b> PD-L1+CTx 1L biliary tract cancer TOPAZ-1</p>
Removed from Phase I	Removed from Phase II	Removed from Phase III	Removed from registration
	<p><b><u>NME</u></b> <b>Imfinzi + FOLFOX + bevacizumab</b> PD-L1 + chemo + VEGF + multiple novel ONC therapies 1L MSS-CRC COLUMBIA 1</p>	<p><b><u>Life-cycle management</u></b> <b>Imfinzi CALLA</b> PD-L1 mAb locally-advanced cervical cancer</p>	<p><b><u>Life-cycle management</u></b> <b>Lynparza<sup>#</sup> OlympiA</b> PARP inhibitor gBRCA adjuvant breast cancer</p>



# Q1 2022 Oncology new molecular entity<sup>1</sup> pipeline

Phase 1 14 New Molecular Entities		Phase 2 16 New Molecular Entities		Phase 3 15 New Molecular Entities	
AZD0466# BCL2/xL haematological malignancies	AZD1390 ATM glioblastoma	adavosertib# Wee1 ovarian / uterine serous / solid tumours	AZD0171 + <i>Imfinzi</i> + CTx anti-LIF+PD-L1+CTx 1L metastatic pancreatic ductal adenocarcinoma	camizestrant + CDK4/6i SERENA-6 SERD+CDK4/6 1L HR+ HER2- ESR1m breast cancer	camizestrant + palbociclib SERENA-4 SERD+CDK4/6 1L HR+ HER2- breast cancer
AZD2936# PD1/TIGIT bispecific mAb solid tumours	AZD5991 MCL1 haematological malignancies	AZD4573 CDK9 haematological malignancies	AZD4573 + <i>Calquence</i> CDK9+BTK haematological malignancies	capivasertib + abiraterone CAPItello-281# AKT+abiraterone PTEN deficient mHSPC	capivasertib + CTx CAPItello-290# AKT+chemotherapy mTNBC 1L
AZD7648# DNAPK solid tumours	AZD7789 PD1/TIM3 bispecific mAb solid tumours, haem malignancies	AZD5305 PARP1Sel solid tumours	camizestrant SERD ER+ breast	capivasertib + fulvestrant + palbociclib CAPItello-292# AKT+fulvestrant+CDK4/6 HR+ breast 1L	capivasertib + fulvestrant CAPItello-291# AKT+fulvestrant locally-advanced (inoperable) or metastatic breast
AZD8205 B7-H4 TOP1i ADC solid tumours	AZD8701 +/- <i>Imfinzi</i> # FOXP3 +/- PD-L1 solid tumours	capivasertib# AKT inhibitor prostate cancer	ceralasertib ATR solid tumours	datopotamab deruxtecan TROPION-Breast01# TROP2 2-3L HR+ HER2- BC	datopotamab deruxtecan TROPION-Lung01# TROP2 2L+ NSCLC without actionable mutation
<i>Imfinzi</i> + adavosertib# PD-L1+Wee1 solid tumours	IPH5201# CD39 solid tumours	<i>Imfinzi</i> (platform) COAST# PD-L1+multiple novel ONC therapies NSCLC	<i>Imfinzi</i> (platform) HUDSON PD-L1+multiple novel ONC therapies post IO NSCLC	<i>Imfinzi</i> + treme + TACE +/- Lenvatinib EMERALD-3 PD-L1+CTLA4+VEGF+/-chemo-embolization LR HCC	<i>Imfinzi</i> + tremelimumab + SoC NILE PD-L1+CTLA-4+SoC 1L urothelial cancer
MEDI1191# IL12 mRNA solid tumours	MEDI5752 + Lenvatinib PD-1/CTLA-4+VEGF advanced renal cell carcinoma	<i>Imfinzi</i> (platform) NeoCOAST# PD-L1+multiple novel ONC therapies NSCLC	<i>Imfinzi</i> + monalizumab# PD-L1+NKG2a solid tumours	Lynparza + <i>Imfinzi</i> + bevacizumab DUO-O# PARP+PD-L1+VEGF 1L ovarian	<i>Imfinzi</i> +/- tremelimumab + CRT ADRIATIC# PD-L1+/-CTLA-4+CRT 1L SCLC
MEDI9253 rNDV IL12 solid tumours	<i>Tagrisso</i> + (Koselugo or savolitinib) TATTON# EGFR+MEK/MET advanced EGFRm NSCLC	<i>Imfinzi</i> + Lynparza ORION# PD-L1+PARP 1L mNSCLC	oleclumab + CTx or <i>Imfinzi</i> + oleclumab + CTx CD73 + chemo or PD-L1 + CD73 + chemo pancreatic	Lynparza + <i>Imfinzi</i> DUO-E# PARP+PD-L1 1L endometrial cancer	monalizumab + cetuximab INTERLINK-1# NKG2a+EGFR 2L+ relapsed metastatic HNSCC
		MEDI5752 PD-1/CTLA-4 solid tumours	<i>Tagrisso</i> + savolitinib SAVANNAH# EGFR+MET advanced EGFRm NSCLC	savolitinib + <i>Imfinzi</i> SAMETA# MET+PD-L1 1L papillary renal cell carcinoma	
<b>Under Review</b>					
2 New Molecular Entity					
			<i>Imfinzi</i> +/- tremelimumab + CTx POSEIDON PD-L1+/-CTLA-4+CTx 1L NSCLC		<i>Imfinzi</i> + tremelimumab HIMALAYA# PD-L1+CTLA-4 1L HCC

Phase progressions based on first patient dose achievement.

<sup>1</sup> Includes novel combinations and additional indications for assets where the lead is not yet launched.

# Partnered and/or in collaboration; ¶ Registrational Phase II/III trial

● Precision medicine approach being explored



# Q1 2022 Oncology life-cycle management<sup>1</sup> pipeline

Phase 1 1 Project	Phase 2 7 Projects	Phase 3 32 Projects			Under Review 4 Projects
<i>Enhertu</i> (platform) DESTINY-Breast08# ADC breast	<i>Enhertu</i> (platform) DESTINY-Breast07# ADC breast	<i>Calquence</i> + R-CHOP ESCALADE BTK+R-CHOP 1L DLBCL	<i>Calquence</i> + venetoclax + obinutuzumab AMPLIFY# BTK+BCL-2+anti-CD20 1st line CLL	<i>Calquence</i> ECHO# BTK inhibitor 1st line MCL	<i>Enhertu</i> DESTINY-Breast03# ADC breast
	<i>Enhertu</i> DESTINY-PanTumour01# HER2 targeting ADC HER2 mutant tumours	<i>Enhertu</i> DESTINY-Breast04# ADC breast	<i>Enhertu</i> DESTINY-Breast02# ADC breast	<i>Enhertu</i> DESTINY-Breast05# ADC breast	<i>Enhertu</i> DESTINY-Lung01# ADC NSCLC
	<i>Enhertu</i> DESTINY-PanTumour02# HER2 targeting ADC HER2-expressing solid tumours	<i>Enhertu</i> DESTINY-Breast06# ADC breast	<i>Enhertu</i> DESTINY-Breast09# ADC breast	<i>Enhertu</i> DESTINY-Breast11# ADC breast	<i>Lynparza</i> + abiraterone PROpel# PARP+NHA prostate cancer
	<i>Imfinzi</i> (platform) BEGONIA PD-L1 1L mTNBC	<i>Enhertu</i> DESTINY-Lung04# ADC NSCLC	<i>Enhertu</i> DESTINY-Gastric04# HER2 targeting ADC HER2+ gastric cancer 2L	<i>Imfinzi</i> PEARL PD-L1 1L metastatic NSCLC	<i>Imfinzi</i> + CTx TOPAZ-1# PD-L1+CTx 1L biliary tract cancer
	<i>Imfinzi</i> (platform) MAGELLAN# PD-L1 1L mNSCLC	<i>Imfinzi</i> + CRT KUNLUN# PD-L1+CRT locally-advanced ESCC	<i>Imfinzi</i> + CRT PACIFIC-2# PD-L1+CRT locally-advanced stage III NSCLC	<i>Imfinzi</i> + CRT PACIFIC-5 (China)# PD-L1+CRT locally-advanced stage III NSCLC	
	<i>Lynparza</i> (basket) MK-7339-002 / LYNK002# PARP inhibitor HRRm cancer	<i>Imfinzi</i> + CTx neoadjuvant AEGEAN# PD-L1+CTx locally-advanced stage II-III NSCLC	<i>Imfinzi</i> + CTx MERMAID-1 PD-L1 stage II-III adjuvant NSCLC	<i>Imfinzi</i> + CTx NIAGARA PD-L1 mAb + CTx muscle invasive bladder cancer	
	<i>Tagrisso</i> ORCHARD platform study# EGFR+multiple novel ONC therapies EGFRm NSCLC	<i>Imfinzi</i> + VEGF + TACE EMERALD-1# PD-L1+VEGF+TACE locoregional HCC	<i>Imfinzi</i> + domvanalimab (AB154) + CTx PACIFIC-8# PD-L1+TIGIT+CTx unresectable stage III NSCLC	<i>Imfinzi</i> +- oleclumab +- mona PACIFIC-9# PD-L1 + NKG2a or PD-L1 + CD73 Unresectable stage III NSCLC	
		<i>Imfinzi</i> MERMAID-2 PD-L1 stage II-III premetastatic NSCLC	<i>Imfinzi</i> + FLOT MATTERHORN# PD-L1+CTx neo-adjuvant/adjuvant gastric cancer	<i>Imfinzi</i> + EV +/- treme VOLGA PD-L1 + nectin-4 targeting ADC +/- CTLA4 muscle invasive bladder cancer	
		<i>Lynparza</i> MONO-OLA1# PARP 1L BRCAwt ovarian cancer	<i>Imfinzi</i> post-SBRT PACIFIC-4# PD-L1 post-SBRT stage I/II NSCLC	<i>Imfinzi</i> + VEGF EMERALD-2# PD-L1+VEGF adjuvant HCC	
		<i>Tagrisso</i> +/- CTx neoadjuvant NeoADAURA EGFR inhibitor +/- CTx stage II/III resectable EGFRm NSCLC	<i>Lynparza</i> LYNK-003# PARP platinum sensitive 1L colorectal cancer	<i>Imfinzi</i> POTOMAC PD-L1 mAb non muscle invasive bladder cancer	
		<i>Tagrisso</i> LAURA EGFR inhibitor stage III EGFRm non-small cell lung cancer	<i>Tagrisso</i> + CTx FLAURA2 EGFR+chemo 1L adv EGFRm NSCLC		

Phase progressions based on first patient dose achievement.

<sup>1</sup> Includes significant life-cycle management projects and parallel indications for assets beyond Phase III

# Partnered and/or in collaboration; \* Registrational Phase II/III trial

● Precision medicine approach being explored



# Q1 2022 BioPharmaceuticals new molecular entity<sup>1</sup> pipeline

Phase 1 14 New Molecular Entities		Phase 2 17 New Molecular Entities		Phase 3 5 New Molecular Entities	Under Review 1 New Molecular Entity
AZD2373 Podocyte health nephropathy	AZD2693 nonalcoholic steatohepatitis	AZD1402# Inhaled IL-4Ra asthma	AZD4831 MPO HFpEF	brazikumab INTREPID IL-23 mAb Crohn's disease	nirsevimab# RSV mAb-YTE passive RSV immunisation
AZD3366 CD39L3 CV disease	AZD3427 Relaxin mimetic CV disease	AZD5718 FLAP coronary artery disease / CKD	AZD7986# DPP1 COPD	eplontersen# LICA ATTR-Cardiomyopathy	
AZD4041# orexin 1 receptor antagonist opioid use disorder	AZD4604 inhaled JAK1 asthma	AZD8233 PCSK9-ASO hypercholesterolemia	AZD8601# VEGF-A cardiovascular disease	eplontersen# LICA hATTR-Polyneuropathy	
AZD5055 Porcupine inhibitor IPF	AZD5462 Relaxin mimetic CV disease	AZD9977 + <i>Farxiga/Forxiga</i> MR+SGLT2 HF with CKD	brazikumab EXPEDITION IL-23 mAb ulcerative colitis	PT027# ICS/SABA asthma	
AZD7503 ASO NASH	AZD8630# Inhaled TSLP Fab asthma	cotadutide GLP-1/glucagon T2D / obesity / NASH / DKD	MEDI6570 LOX-1 CV disease	tozorakimab IL-33 mAb COPD	
MEDI0618 PAR2 antagonist mAb osteoarthritis pain	MEDI1341# alpha synuclein mAb parkinson's disease	MEDI7352 NGF/TNF OA pain / painful diabetic neuropathy	navafenterol# MABA chronic obstructive pulmonary disease		
MEDI1814# amyloid beta mAb alzheimer's disease	MEDI8367 avb8 chronic kidney disease	tozorakimab IL-33 mAb diabetic kidney disease	tozorakimab IL-33 mAb asthma		
		tozorakimab IL-33 mAb COVID-19	tozorakimab IL-33 mAb atopic dermatitis		
		zibotentan + <i>Farxiga/Forxiga</i> ZENITH-CKD ETA antagonist+SGLT2 CKD			

Phase progressions based on first patient dose achievement.

<sup>1</sup> Includes novel combinations and additional indications for assets where the lead is not yet launched

# Partnered and/or in collaboration; ¶ Registrational Phase II/III trial

● Precision medicine approach being explored



# Q1 2022 BioPharmaceuticals life-cycle management<sup>1</sup> pipeline

Phase 1 0 Projects	Phase 2 6 Projects	Phase 3 16 Projects	Under Review 0 Projects
	<i>Andexxa</i> (ALXN2070) anti-factor Xa reversal urgent surgery	<i>Breztri/Trixeo</i> (PT010) KALOS LOGOS LABA/LAMA/ICS asthma	<i>Farxiga/Forxiga</i> DAPA-MI SGLT2 prevention of HF and CV death following a myocardial infarction
	<i>Fasenra</i> ARROYO IL-5R mAb chronic spontaneous urticaria	<i>Farxiga/Forxiga</i> DELIVER SGLT2 HFpEF	<i>Fasenra</i> FJORD IL-5R mAb bullous pemphigoid
	<i>Fasenra</i> HILLIER IL-5R mAb atopic dermatitis	<i>Fasenra</i> HUDSON IL-5R mAb eosinophilic gastritis and eosinophilic gastroenteritis	<i>Fasenra</i> MAHALE IL-5R mAb non-cystic fibrosis bronchiectasis
	roxadustat# hypoxia-inducible factor prolyl hydroxylase inhibitor chemotherapy induced anaemia	<i>Fasenra</i> MANDARA IL-5R EGPA	<i>Fasenra</i> MESSINA IL-5R mAb eosinophilic esophagitis
	<i>Saphnelo</i> # Type I IFN receptor mAb lupus nephritis	<i>Fasenra</i> NATRON IL-5R mAb hypereosinophilic syndrome	<i>Fasenra</i> ORCHID# IL-5R mAb nasal polyps
	<i>Tezspire</i> COURSE# TSLP mAb chronic obstructive pulmonary disease	<i>Fasenra</i> RESOLUTE# IL-5R COPD	<i>Lokelma</i> DIALIZE-Outcomes potassium binder CV outcomes in patients on chronic haemodialysis with hyperkalaemia
		<i>Lokelma</i> STABILIZE-CKD potassium binder hyperkalaemia in CKD	roxadustat# HIFPH anaemia MDS
		<i>Saphnelo</i> TULIP-SC# Type I IFN receptor SLE SC	<i>Tezspire</i> WAYPOINT# TSLP mAb nasal polyps

Phase progressions based on first patient dose achievement.

<sup>1</sup> Includes significant life-cycle management projects and parallel indications for assets beyond Phase III

# Partnered and/or in collaboration; ¶ Registrational Phase II/III trial

● Precision medicine approach being explored



# Q1 2022 Rare Disease pipeline<sup>1</sup>

Phase 1 4 Projects	Phase 2 3 Projects	Phase 3 7 Projects	Under Review 2 Projects
ALXN1720 anti-C5 bi-specific minibody generalised myasthenia gravis	ALXN2050 oral factor D inhibitor paroxysmal nocturnal haemoglobinuria	acoramidis (ALXN2060)# oral TTR stabilizer transthyretin amyloid cardiomyopathy	<i>Ultomiris</i> (ALXN1210) anti-complement C5 mAb generalised myasthenia gravis
ALXN1820 anti-properdin bi-specific haematology	danicopan (ALXN2040) factor D inhibitor geographic atrophy	ALXN1840 bis-choline tetrathiomolybdate Wilsons disease	<i>Ultomiris</i> (ALXN1210) anti-complement C5 mAb subcutaneous, paroxysmal nocturnal haemoglobinuria and atypical haemolytic uraemic syndrome
ALXN1850 next generation asfotase alfa hypophosphatasia	<i>Ultomiris</i> (ALXN1210) anti-complement C5 mAb dermatomyositis	CAEL-101 fibril-reactive mAb AL amyloidosis	
NI006# TTR depleter transthyretin amyloid cardiomyopathy		danicopan (ALXN2040) factor D inhibitor paroxysmal nocturnal haemoglobinuria with extravascular haemolysis	
		<i>Ultomiris</i> (ALXN1210) anti-complement C5 mAb neuromyelitis optica spectrum disorder	
		<i>Ultomiris</i> (ALXN1210) anti-complement C5 mAb haematopoietic stem cell transplant-associated thrombotic microangiopathy	
		<i>Ultomiris</i> (ALXN1210) anti-complement C5 mAb complement-mediated thrombotic microangiopathy	

Phase progressions based on first patient dose achievement.

<sup>1</sup> Includes new molecular entities and significant life-cycle management projects

# Partnered and/or in collaboration; ¶ Registrational Phase II/III trial

● Precision medicine approach being explored



# Estimated key regulatory submission acceptances

	NME		LCM	
	PT027 asthma	eplontersen hATTR-PN	danicopan (ALXN2040) PNH-EVH	eplontersen ATTR-CM
	H1 2022	H2 2022	ALXN1840 Wilson's disease	ATTR-CM
	Enhertu DESTINY-Breast04	Enhertu DESTINY-Breast02	Imfinzi +/- tremelimumab + CRT LDS-SCLC ADRIATIC	Imfinzi + treme + TACE +/- Lenvatinib EMERALD-3
		Imfinzi + VEGF + TACE locoregional HCC EMERALD-1	Imfinzi + tremelimumab + SoC urothelial NILE	Orpathys + Imfinzi 1L PRCC SAMETA
		Farxiga/Forxiga (HFpEF) DELIVER	datopotamab deruxtecan NSCLC TROPION-Lung01	monalizumab + cetuximab INTERLINK-1
		Ultomiris (ALXN1210) neuromyelitis optica spectrum disorder	capivasertib + fulvestrant 2L locally advanced or mBC CAPitello-291	Lynparza + Imfinzi endometrial cancer DUO-E
			capivasertib + CTx 1L mTNBC CAPitello-290	Lynparza + Imfinzi+ bevacizumab ovarian DUO-O
			<b>2023</b>	<b>&gt;2023</b>
			Imfinzi NSCLC PEARL	Imfinzi +CTx stage II-III adjuvant NSCLC MERMAID-1
			Imfinzi + CRT NSCLC PACIFIC-5 (China)	Imfinzi + EV +/- treme VOLGA
			Imfinzi + CRT NSCLC PACIFIC-2	Imfinzi + CRT neo-adj/adj gastric MATTERHORN
			Imfinzi + CTx neoadjuvant NSCLC AEGEAN	Imfinzi post-SBRT NSCLC PACIFIC-4
			Imfinzi + CTx muscle invasive bladder NIAGARA	Imfinzi stage II-III premetastatic NSCLC MERMAID-2
			Imfinzi + VEGF adjuvant HCC EMERALD-2	Imfinzi non muscle invasive bladder POTOMAC
			Tagrisso locally adv. unresectable NSCLC LAURA	Lynparza platinum sensitive 1L colorectal LYNK-003
			Tagrisso + CTx EGFRm NSCLC FLAURA2	Lynparza 1L BRCAwt ovarian MONO-OLA1
			Fasenna eosinophilic esophagitis MESSINA	Tagrisso stage II-III resectable EGFRm NSCLC NeoADAURA
				Imfinzi + domvanalimab (AB154) + CTx PACIFIC-8
				Imfinzi +- oleclumab +- mona unresectable stage III NSCLC PACIFIC-9
				Fasenna nasal polyps ORCHID
				Fasenna EG/EGE HUDSON
				CAEL-101 AL amyloidosis
				acoramidis (ALXN2060) transthyretin amyloid cardiomyopathy
				brazikumab crohns disease INTREPID
				Farxiga prevention of HF and CV death following myocardial infarction DAPA-MI
				Lokelma DIALIZE-Outcomes
				Lokelma STABILIZE-CKD
				roxadustat anemia in MDS
				Bretri/Trixio asthma KALOS, LOGOS
				Fasenna COPD RESOLUTE
				Fasenna bullous pemphigoid FJORD
				Fasenna non-cystic fibrosis bronchiectasis MAHALE
				Fasenna EGPA MANDARA
				Fasenna HES NATRON
				Ultomiris dermatomyositis
				Ultomiris haematopoietic STC-associated TMA
				Ultomiris complement-mediated TMA



# Designations in our pipeline

3

Accelerated approvals

Andexxa Acute Major Bleed (US)
Calquence MCL 1L (US)
nirsevimab RSV mAb-YTE (EU)

18

Breakthrough / PRIME<sup>1</sup> / Sakigake<sup>2</sup>

danicopan (ALXN2040) PNH-EVH (US)
danicopan (ALXN2040) PNH-EVH (EU)
Calquence CLL ELEVATE-TN, ASCEND (US)
Calquence MCL 1L (US)
Enhertu HER2+ breast 2L DESTINY-Breast03 (US)
Enhertu HER2+ breast 2L DESTINY-Breast03 (CN)
Enhertu HER2-low, unresectable and/or metastatic breast cancer DESTINY-Breast04 (US)
Enhertu HER2+/HER2low gastric 3L DESTINY-Gastric01 (US)
Enhertu HER2+/HER2low gastric 3L DESTINY-Gastric01 (JP) <sup>2</sup>
Enhertu HER2mut NSCLC 2L+ DESTINY-Lung01 (US)
Forxiga CKD DAPA-CKD (US)
Koselugo NFI type 1 SPRINT (US)
nirsevimab RSV mAb-YTE MELODY-MEDLEY (US)
nirsevimab RSV mAb-YTE MELODY-MEDLEY (CN)
nirsevimab RSV mAb-YTE MELODY-MEDLEY (EU) <sup>1</sup>
Soliris GBS (JP) <sup>2</sup>
Tagrisso adjuvant NSCLC ADAURA (US)
Tezspire asthma NAVIGATOR (US)

8

Fast Track

Saphnelo SLE (US)
CAEL-101 AL amyloidosis (US)
cotadutide NASH (US)
Fasenra EG/EGE HUDSON (US)
Forxiga CKD DAPA-CKD (US)
Forxiga MI RRCT DAPA-MI (US)
Lokelma ESRD DIALIZE-OUTCOMES (US)
nirsevimab RSV mAb-YTE MELODY-MEDLEY (US)

14

Priority Review

Brilinta stroke THALES (US)
Calquence MCL 1L (US)
Enhertu HER2+/HER2low gastric 3L DESTINY-Gastric01 (US)
Enhertu HER2+ breast 2L DESTINY-Breast03 (US)
Enhertu HER2mut NSCLC 2L+ DESTINY-Lung01 (US)
Forxiga CKD DAPA-CKD (US)
Forxiga CKD DAPA-CKD (JP)
Imfinzi + tremelimumab HCC HIMALAYA (US)
Koselugo NFI type 1 SPRINT (US)
Lynparza gBRCA adj breast OlympiA (US)
Roxadustat chronic kidney disease (CN)
Tagrisso adjuvant NSCLC (ADAURA) (US)
Tezspire asthma NAVIGATOR (US)
Ultomiris gMG (US)

30

Orphan

danicopan (ALXN2040) PNH (US)
danicopan (ALXN2040) PNH (EU)
ALXN2050 PNH (US)
ALXN2050 PNH (EU)
ALXN1840 WD (US)
ALXN1840 WD (EU)
Andexxa Acute Major Bleed (JP)
CAEL-101 AL amyloidosis (US)
CAEL-101 AL amyloidosis (EU)
Calquence CLL 1L (US)
Calquence CLL 1L (EU)
Calquence MCL 1L (US)
Enhertu HER2+/HER2low gastric 3L DESTINY-Gastric01 (US)
eplontersen transthyretin-mediated amyloidosis (US)
Fasenra EG/EGE HUDSON (US)
Fasenra EGPA MANDARA (US)
Fasenra EOE MESSINA (US)
Fasenra HES NATRON (US)
Imfinzi + CTx Biliary Tract 1L TOPAZ-1 (US)
Imfinzi + CTx Biliary Tract 1L TOPAZ-1 (JP)
Imfinzi +/- treme HCC 1L (EU)
Imfinzi +/- treme HCC 1L (US)
Koselugo NFI type 1 SPRINT (US)
Koselugo NFI type 1 SPRINT (EU)
Koselugo NFI type 1 SPRINT (JP)
Lynparza gBRCA adj breast OlympiA (JP)
Tezspire Eosinophilic Esophagitis (US)
Ultomiris DM (US)
Ultomiris HSCT-TMA (US)
Ultomiris SC PNH (US)

ACCELERATED APPROVAL, these regulations allowed medicines for serious conditions that addressed an unmet medical need to be approved based on a surrogate endpoint.

BREAKTHROUGH DESIGNATION is a process designed to expedite the development and review of medicines which may demonstrate substantial improvement over available therapy. <sup>1</sup>PRIME is a scheme launched by the EMA to enhance support for the development of medicines that target an unmet medical need. <sup>2</sup>SAKIGAKE is aimed at early introduction of innovative medicines, medical devices, etc. that are initially developed in Japan.


FAST TRACK is a process designed to facilitate the development, and expedite the review of medicines to treat serious conditions and fill an unmet medical need.

PRIORITY REVIEW DESIGNATION is the US FDA's goal to take action on an application within 6 months.

ORPHAN DRUG DESIGNATION, intended for treatment, diagnosis or prevention of rare diseases/disorders that affect fewer than 200,000 patients in the US, or that affect more than 200,000 patients but are not expected to recover the costs of developing and marketing a treatment drug.



Oncology –  
approved medicines and  
late-stage pipeline



# Tagrisso (highly-selective, irreversible EGFRi)

## NSCLC

Trial	Population	Patients	Design	Endpoints	Status
Phase III ADAURA NCT02511106	Adjuvant EGFRm NSCLC	682	<ul style="list-style-type: none"> <li>Arm 1: <i>Tagrisso</i> QD following complete tumour resection, with or without chemo</li> <li>Arm 2: placebo</li> <li>Global trial - 25 countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: DFS</li> <li>Secondary endpoints: DFS Rate, OS, OS Rate, QoL</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2015</li> <li>LPCD: Q1 2019</li> <li>Data readout: Q2 2020</li> <li>Trial unblinded due to efficacy</li> <li>DFS primary endpoint met</li> </ul>
Phase III LAURA NCT03521154	Maintenance therapy in patients with locally advanced, unresectable EGFRm Stage III NSCLC whose disease has not progressed following platinum-based chemoradiation therapy	200	<ul style="list-style-type: none"> <li>Arm 1: <i>Tagrisso</i></li> <li>Arm 2: placebo</li> <li>Global trial - 17 countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: PFS (BICR)</li> <li>Secondary endpoints: CNS PFS, OS, DoR, ORR, DCR</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2018</li> <li>Data anticipated: 2023</li> </ul>
Phase II ELIOS NCT03239340	EGFR TKI treatment-naïve patients with locally-advanced or metastatic EGFRm NSCLC	150	<ul style="list-style-type: none"> <li>Single arm trial - <i>Tagrisso</i></li> <li>Global trial - five countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: proportion of patients with a given tumour genetic and proteomic marker at the point of disease progression as defined by the investigator</li> <li>Secondary endpoints: PFS, ORR, DoR</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2018</li> </ul>
Phase I ODIN-BM NCT03463525	Patients with EGFRm NSCLC with brain metastases	8	<ul style="list-style-type: none"> <li>Single-arm trial - <i>Tagrisso</i></li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: assessments of brain standard uptake value (SUV) and pharmacokinetics (PK)</li> <li>Secondary endpoint: PK</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2018</li> <li>LPCD: Q1 2020</li> <li>Data readout: Q4 2021</li> </ul>



# Tagrisso (highly-selective, irreversible EGFRi)

## NSCLC, combinations

Trial	Population	Patients	Design	Endpoints	Status
Phase III NeoADAURA NCT04351555	Neoadjuvant EGFRm NSCLC	351	<ul style="list-style-type: none"> <li>Arm 1: placebo plus pemetrexed/carboplatin or pemetrexed/cisplatin</li> <li>Arm 2: <i>Tagrisso</i> plus pemetrexed/carboplatin or pemetrexed/cisplatin</li> <li>Arm 3: <i>Tagrisso</i></li> <li>Global trial – 23 countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: mPR</li> <li>Secondary endpoints: cPR, EFS, DFS, OS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2021</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase III FLAURA2 NCT04035486	1st-line EGFRm NSCLC	586	<ul style="list-style-type: none"> <li>Arm 1: <i>Tagrisso</i> plus pemetrexed/carboplatin or pemetrexed/cisplatin</li> <li>Arm 2: <i>Tagrisso</i></li> <li>Global trial – 23 countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: PFS</li> <li>Secondary endpoints: OS, LOS, ORR DoR, Depth of response, PFS2. QoL, PK</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2019</li> <li>Data anticipated: 2023</li> </ul>
Phase III COMPEL NCT04765059	EGFRm metastatic NSCLC patients who have progressed extracranially following 1L treatment with <i>Tagrisso</i>	204	<ul style="list-style-type: none"> <li>Arm 1: <i>Tagrisso</i> plus pemetrexed/carboplatin or pemetrexed/cisplatin</li> <li>Arm 2: placebo plus pemetrexed/carboplatin or pemetrexed/cisplatin</li> <li>Global trial</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: PFS</li> <li>Secondary endpoints: Intracranial PFS, Extracranial PFS, OS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2021</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase III SAFFRON NCT05261399 Partnered	EGFR mutated, MET-overexpressed and/or amplified, locally advanced or metastatic NSCLC patients who have progressed on first- or second-line treatment with <i>Tagrisso</i>	324	<ul style="list-style-type: none"> <li>Arm 1: <i>Tagrisso</i> + <i>Orpathys</i></li> <li>Arm2: pemetrexed with either cisplatin or carboplatin</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: PFS</li> <li>Secondary endpoints: OS, ORR, PK, DCR, DoR</li> </ul>	<ul style="list-style-type: none"> <li>Initiating</li> </ul>
Phase II SAVANNAH NCT03778229	EGFRm / MET+, locally advanced or metastatic NSCLC who have progressed following treatment with <i>Tagrisso</i>	259	<ul style="list-style-type: none"> <li>Single arm trial: <i>Tagrisso</i> + <i>Orpathys</i></li> <li>Global trial</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: ORR</li> <li>Secondary endpoints include PFS, DoR and OS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2019</li> <li>Data anticipated: H2 2022</li> </ul>



# Tagrisso (highly-selective, irreversible EGFRi)

## NSCLC, combinations

Trial	Population	Patients	Design	Endpoints	Status
Phase II ORCHARD NCT03944772	Advanced EGFRm NSCLC patients who have progressed on first line Tagrisso treatment	182	<ul style="list-style-type: none"> <li>Modular design platform trial:</li> <li>Module 1: <i>Tagrisso</i> + <i>Orpathys</i> (cMET)</li> <li>Module 2: <i>Tagrisso</i> + gefitinib (EGFRm)</li> <li>Module 3: <i>Tagrisso</i> + necitumumab (EGFRm)</li> <li>Module 4: carboplatin + pemetrexed + <i>Imfinzi</i></li> <li>Module 5: <i>Tagrisso</i> + alectinib (ALK)</li> <li>Module 6: <i>Tagrisso</i> + selpercatinib (RET fusion)</li> <li>Module 7: <i>Imfinzi</i> + etoposide + carboplatin or cisplatin</li> <li>Module 8: <i>Tagrisso</i> + pemetrexed + carboplatin or cisplatin</li> <li>Module 9: <i>Tagrisso</i> + <i>Koselugo</i></li> <li>No intervention: observational cohort</li> <li>Global trial - 8 countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: ORR</li> <li>Secondary endpoints: PFS, DoR, OS, safety and tolerability</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2019</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase II NCT04606771	EGFRm/MET amplified advanced NSCLC	56	<ul style="list-style-type: none"> <li><i>Tagrisso</i> and <i>Orpathys</i> contribution of components</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: ORR</li> <li>Secondary endpoints: PFS, DoR, OS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2020</li> <li>Data anticipated: H2 2022</li> </ul>



# Imfinzi (PD-L1 mAb)

## Lung cancer, early disease

Trial	Population	Patients	Design	Endpoints	Status
Phase II HUDSON NCT03334617	NSCLC, patients who progressed on an anti-PD-1/PD-L1 containing therapy	340	<ul style="list-style-type: none"> <li>Open-label, biomarker-directed, multicentre trial</li> <li>Module 1: <i>Imfinzi</i> and <i>Lynparza</i></li> <li>Module 2: <i>Imfinzi</i> and danvatirsen</li> <li>Module 3: <i>Imfinzi</i> and ceralasertib (AZD6738)</li> <li>Module 4: <i>Imfinzi</i> and vistusertib</li> <li>Module 5: <i>Imfinzi</i> and oleclumab</li> <li>Module 6: <i>Imfinzi</i> and <i>Enhertu</i></li> <li>Module 7: <i>Imfinzi</i> and cediranib</li> <li>Module 8: ceralasertib</li> <li>Module 9: <i>Imfinzi</i> and ceralasertib</li> </ul>	<ul style="list-style-type: none"> <li>Primary outcome: ORR</li> <li>Secondary outcomes: efficacy including OS, PFS, DCR, and safety and tolerability, DoR</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2018</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase II COAST NCT03822351	Stage III NSCLC unresectable	189	<ul style="list-style-type: none"> <li>Arm A: <i>Imfinzi</i></li> <li>Arm B: <i>Imfinzi</i> + oleclumab</li> <li>Arm C: <i>Imfinzi</i> + monalizumab</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: OR per RECIST v1.1</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2018</li> <li>Data readout: Q3 2021</li> </ul>
Phase II NeoCOAST NCT03794544	Resectable, early-stage NSCLC	84	<ul style="list-style-type: none"> <li>Arm A: <i>Imfinzi</i></li> <li>Arm B: <i>Imfinzi</i> + oleclumab</li> <li>Arm C: <i>Imfinzi</i> + monalizumab</li> <li>Arm D: <i>Imfinzi</i> + danvatirsen</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: Major pathological response rate</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2019</li> <li>Data readout: Q1 2022</li> </ul>
Phase I/II SCoPe-D1 NCT04870112	NSCLC SCLC	124	<ul style="list-style-type: none"> <li>Open-label, multicentre trial to evaluate the safety, pharmacokinetics, and preliminary efficacy of subcutaneous <i>Imfinzi</i></li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: PK, safety</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2021</li> <li>Data anticipated: &gt;2023</li> </ul>



# Imfinzi (PD-L1 mAb)

## NSCLC, early disease

Trial	Population	Patients	Design	Endpoints	Status
Phase III MERMAID-1 NCT04385368	Completely resected Stage II and III NSCLC	332	<ul style="list-style-type: none"> <li>Arm 1: <i>Imfinzi</i> + SoC chemo</li> <li>Arm 2: placebo + SoC chemo</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: DFS</li> <li>Secondary endpoints: DFS, OS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2020</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase III MERMAID-2 NCT04642469	Completely resected Stage II-III NSCLC	284	<ul style="list-style-type: none"> <li>Arm 1: <i>Imfinzi</i></li> <li>Arm 2: placebo</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: DFS</li> <li>Secondary endpoints: DFS, PFS, OS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2021</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase III AEGEAN NCT03800134	Neoadjuvant NSCLC patients Stage II and III resected NSCLC (incl. EGFR/ALK positive)	800	<ul style="list-style-type: none"> <li>Arm 1: <i>Imfinzi</i> + platinum-based chemo</li> <li>Arm 2: placebo + platinum-based chemo</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: pCR, EFS</li> <li>Secondary endpoint: mPR</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2019</li> <li>Data anticipated: 2023</li> </ul>
Phase III ADJUVANT BR.31 NCT02273375 Partnered	Adjuvant NSCLC patients Stage Ib (≥4cm) – stage IIIa resected (incl. EGFR/ALK positive)	1360	<ul style="list-style-type: none"> <li>Arm 1: <i>Imfinzi</i> mg/kg i.v. Q4W x 12m</li> <li>Arm 2: placebo</li> <li>Global trial</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: DFS</li> <li>Secondary endpoint: OS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2015</li> <li>LPCD: Q1 2020</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase III PACIFIC-2 NCT03519971	Unresected, locally-advanced NSCLC	300	<ul style="list-style-type: none"> <li>Arm 1: <i>Imfinzi</i> i.v. Q4W + chemo/RT</li> <li>Arm 2: placebo + chemo/RT</li> <li>ex US global trial</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: PFS</li> <li>Secondary endpoints: OS, ORR</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2018</li> <li>LPCD: Q3 2019</li> <li>Data anticipated: H2 2022</li> </ul>
Phase III PACIFIC-4 NCT03833154	<i>Imfinzi</i> with SBRT in unresected, Stage I/II NSCLC	630	<ul style="list-style-type: none"> <li>Arm 1: <i>Imfinzi</i> i.v. Q4W with definitive SBRT</li> <li>Arm 2: placebo with definitive SBRT</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: PFS</li> <li>Secondary endpoint: OS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2019</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase III PACIFIC-5 NCT03706690	Unresected, locally-advanced NSCLC	360	<ul style="list-style-type: none"> <li>Arm 1: <i>Imfinzi</i> i.v. Q4W following chemo/RT</li> <li>Arm 2: placebo following chemo/RT</li> <li>ex US global trial, China focus</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: PFS</li> <li>Secondary endpoint: OS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2019</li> <li>Data anticipated: 2023</li> </ul>
Phase III PACIFIC-8 NCT05211895 Partnered	Unresected, locally-advanced NSCLC	860	<ul style="list-style-type: none"> <li>Arm 1: <i>Imfinzi</i> + domvanlimab following chemo/RT</li> <li>Arm 2: <i>Imfinzi</i> + PBO following chemo/RT</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: PFS</li> <li>Secondary endpoint: OS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2022</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase III PACIFIC-9 NCT05221840 Partnered	Patients with locally advanced (Stage III), unresectable NSCLC, who have not progressed following platinum-based cCRT	999	<ul style="list-style-type: none"> <li>Arm 1: <i>Imfinzi</i> + oleclumab</li> <li>Arm 2: <i>Imfinzi</i> + monalizumab + placebo</li> <li>Arm 3: <i>Imfinzi</i> + placebo</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: PFS</li> <li>Secondary endpoint: OS, ORR, DoR, PFS2, TFST</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2022</li> <li>Data anticipated: &gt;2023</li> </ul>



# Imfinzi (PD-L1 mAb)

## Other cancers, advanced disease

Trial	Population	Patients	Design	Endpoints	Status
Phase III TOPAZ-1 NCT03875235	BTC 1L	810	<ul style="list-style-type: none"> <li>Arm 1: <i>Imfinzi</i> + gemcitabine + cisplatin</li> <li>Arm 2: placebo + gemcitabine + cisplatin</li> <li>Global trial</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: OS</li> <li>Secondary endpoints: PFS, ORR, DoR</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2019</li> <li>LPCD: Q4 2020</li> <li>Data readout: Q4 2021</li> <li>Primary endpoint met</li> </ul>



# Imfinzi (PD-L1 mAb)

## Other cancers

Trial	Population	Patients	Design	Endpoints	Status
Phase III POTOMAC NCT03528694	Non-muscle invasive bladder cancer	1018	<ul style="list-style-type: none"> <li>Arm 1: BCG (Induction + maintenance)</li> <li>Arm 2: <i>Imfinzi</i> + BCG (Induction only)</li> <li>Arm 3: <i>Imfinzi</i> + BCG (Induction + maintenance)</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: DFS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2018</li> <li>LPCD: Q4 2020</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase III NIAGARA NCT03732677	Muscle-invasive bladder cancer	988	<ul style="list-style-type: none"> <li>Arm 1: <i>Imfinzi</i> in combination with gemcitabine + cisplatin, <i>Imfinzi</i> maintenance</li> <li>Arm 2: gemcitabine + cisplatin</li> </ul>	<ul style="list-style-type: none"> <li>Coprimary endpoints: pCR, EFS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2018</li> <li>LPCD: Q3 2021</li> <li>Data anticipated: 2023</li> </ul>
Phase III EMERALD-1 NCT03778957	Locoregional HCC	710	<ul style="list-style-type: none"> <li>Arm 1: TACE in combination with <i>Imfinzi</i></li> <li>Arm 2: TACE in combination with <i>Imfinzi</i> + bevacizumab</li> <li>Arm 3: TACE in combination with placebo</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: PFS for Arm 2 vs Arm 3</li> <li>Secondary endpoint: PFS for Arm 1 vs Arm 3, OS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2019</li> <li>LPCD: Q3 2021</li> <li>Data anticipated: H2 2022</li> </ul>
Phase III EMERALD-2 NCT03847428	Adjuvant therapy in HCC	888	<ul style="list-style-type: none"> <li>Arm 1: <i>Imfinzi</i> + bevacizumab</li> <li>Arm 2: <i>Imfinzi</i> + placebo</li> <li>Arm 3: placebo + placebo</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: RFS for Arm 1 vs Arm 3</li> <li>Secondary endpoints: RFS Arm 2 vs Arm 3, OS, RFS at 24m</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2019</li> <li>Data anticipated: 2023</li> </ul>
Phase III KUNLUN NCT04550260	Locally advanced, unresectable ESCC	600	<ul style="list-style-type: none"> <li>Arm 1: <i>Imfinzi</i> + definitive CRT</li> <li>Arm 2: placebo + definitive CRT</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: PFS</li> <li>Secondary endpoint: OS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2020</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase III MATTERHORN NCT04592913	Resectable GC/GEJC	900	<ul style="list-style-type: none"> <li>Arm 1: <i>Imfinzi</i> + FLOT</li> <li>Arm 2: placebo + FLOT</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: EFS</li> <li>Secondary endpoints: OS Arm 1 vs Arm 2, pCR Arm 1 vs Arm 2</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2020</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase III SAMETA NCT05043090	MET-Driven, unresectable and locally advanced or metastatic papillary renal cell carcinoma	200	<ul style="list-style-type: none"> <li><i>Orpathys</i> + <i>Imfinzi</i> versus sunitinib and <i>Imfinzi</i> monotherapy</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: PFS</li> <li>Secondary endpoints include OS, ORR, DoR and DCR</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2021</li> <li>Data anticipated: &gt;2023</li> </ul>



# Imfinzi (PD-L1 mAb)

## Other cancers

Approved medicines

Late-stage development

Early development

Trial	Population	Patients	Design	Endpoints	Status
Phase III EMERALD-3 NCT05301842	Locoregional Hepatocellular Carcinoma	525	<ul style="list-style-type: none"><li>• Arm A: TACE + T300 + D + Lenva</li><li>• Arm B: TACE + T300 + D</li><li>• Arm C: TACE</li></ul>	<ul style="list-style-type: none"><li>• Progression Free Survival (PFS) for Arm A vs Arm C</li><li>• Progression Free Survival (PFS) for Arm B vs Arm C</li><li>• Overall Survival (OS) for Arm A vs Arm C</li><li>• Overall Survival (OS) for Arm B vs Arm C</li></ul>	<ul style="list-style-type: none"><li>• FPCD: Q2 2022</li><li>• Data anticipated: &gt;2023</li></ul>

Oncology

CVRM

R&I

Other

V&I

Rare Disease



# Imfinzi (PD-L1 mAb) +/- treme (CTLA-4 mAb)

## Lung cancer, advanced disease

Trial	Population	Patients	Design	Endpoints	Status
Phase III PEARL NCT03003962	NSCLC 1L	650	<ul style="list-style-type: none"> <li>Arm 1: <i>Imfinzi</i> Q4W</li> <li>Arm 2: chemotherapy</li> <li>Asia trial</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: OS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2017</li> <li>LPCD: Q1 2019</li> <li>Data anticipated: H2 2022</li> </ul>
Phase III POSEIDON NCT03164616	NSCLC 1L	1000	<ul style="list-style-type: none"> <li>Arm 1: <i>Imfinzi</i> + chemo</li> <li>Arm 2: <i>Imfinzi</i> + tremelimumab + chemo</li> <li>Arm 3: SoC</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: OS, PFS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2017</li> <li>LPCD: Q4 2018</li> <li>Data readout: Q4 2019</li> <li>PFS primary endpoint met</li> <li>OS data readout Q2 2021</li> </ul>
Phase II MAGELLAN NCT03819465	NSCLC 1L	212	<ul style="list-style-type: none"> <li>Arm A1: <i>Imfinzi</i></li> <li>Arm A2: <i>Imfinzi</i> + danvatirsen</li> <li>Arm A3: <i>Imfinzi</i> + oleclumab</li> <li>Arm A3: MEDI5752</li> <li>Arm B1: <i>Imfinzi</i> + Investigator's choice of chemo</li> <li>Arm B2: <i>Imfinzi</i> + danvatirsen + Investigator's choice of chemo</li> <li>Arm B3: <i>Imfinzi</i> + oleclumab + Investigator's choice of chemo</li> <li>Arm B4: MEDI5752</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: safety &amp; tolerability</li> <li>Secondary endpoints: ORR, DoR, PFS, OS, PK, ADA</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2019</li> <li>Data anticipated: &gt;2023</li> </ul>



# *Imfinzi* (PD-L1 mAb) +/- treme (CTLA-4 mAb)

## Other cancers, advanced disease

Trial	Population	Patients	Design	Endpoints	Status
Phase III NILE NCT03682068	Bladder cancer 1L	1292	<ul style="list-style-type: none"> <li>Arm 1: <i>Imfinzi</i> + tremelimumab + SoC</li> <li>Arm 2: <i>Imfinzi</i> + SoC</li> <li>Arm 3: SoC</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: OS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2018</li> <li>LPD: Q2 2021</li> <li>Data anticipated: 2023</li> </ul>
Phase III HIMALAYA NCT03298451	HCC 1L	1324	<ul style="list-style-type: none"> <li>Arm 1: <i>Imfinzi</i> + tremelimumab</li> <li>Arm 2: <i>Imfinzi</i></li> <li>Arm 3: sorafenib</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: OS</li> <li>Secondary endpoints: PFS, TTP, ORR</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2017</li> <li>LPD: Q4 2019</li> <li>Data readout: Q4 2021</li> <li>Primary endpoint met</li> </ul>
Phase III VOLGA NCT04960709	Muscle invasive bladder cancer ineligible to cisplatin	830	<ul style="list-style-type: none"> <li>Arm 1: <i>Imfinzi</i> + tremelimumab + enfortumab vedotin</li> <li>Arm 2: <i>Imfinzi</i> + enfortumab vedotin</li> <li>Arm 3: SoC cystectomy</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: safety, EFS, pCR</li> <li>Secondary endpoints: OS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2021</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase II BEGONIA NCT03742102	mTNBC 1L	203	<ul style="list-style-type: none"> <li>Arm 1 <i>Imfinzi</i> + paclitaxel</li> <li>Arm 2 <i>Imfinzi</i> + paclitaxel + capivasertib</li> <li>Arm 5 <i>Imfinzi</i> + paclitaxel + oleclumab</li> <li>Arm 6 <i>Imfinzi</i> + <i>Enhertu</i></li> <li>Arm 7 <i>Imfinzi</i> + datopotamab deruxtecan</li> <li>Global trial</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: safety and tolerability</li> <li>Secondary endpoints: ORR, PFS, DoR, OS, PK, ADA</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2019</li> <li>Data anticipated: 2023</li> </ul>
Phase I CLOVER NCT03509012	HNSCC, NSCLC, SCLC	167	<ul style="list-style-type: none"> <li><i>Imfinzi</i> +/- treme in combination with chemoradiation in advanced solid tumours</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: safety</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2018</li> <li>Data readout Q4 2021</li> </ul>



# *Imfinzi* (PD-L1 mAb) +/- treme (CTLA-4 mAb)

## SCLC, advanced disease

Trial	Population	Patients	Design	Endpoints	Status
Phase III ADRIATIC NCT03703297	Limited stage SCLC 1L following platinum-based concurrent chemoradiation therapy	600	<ul style="list-style-type: none"> <li>Arm 1: <i>Imfinzi</i> + tremelimumab (4 doses)</li> <li>Arm 2: <i>Imfinzi</i></li> <li>Arm 3: placebo</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: PFS, OS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2018</li> <li>Data anticipated: H2 2022</li> </ul>



# Lynparza (PARP inhibitor)

## Imfinzi combinations

Trial	Population	Patients	Design	Endpoints	Status
Phase III DUO-O NCT03737643	Advanced ovarian cancer 1L	1256	<ul style="list-style-type: none"> <li>Non tBRCAm (tumour BRCA) patients</li> <li>Arm 1: bevacizumab</li> <li>Arm 2: bevacizumab + <i>Imfinzi</i></li> <li>Arm 3: bevacizumab + <i>Imfinzi</i> + <i>Lynparza</i></li> <li>tBRCAm patients</li> <li>bevacizumab (optional) + <i>Imfinzi</i> + <i>Lynparza</i></li> <li>Global trial</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: PFS</li> <li>Secondary endpoints: OS, PFS2</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2019</li> <li>Data anticipated: 2023</li> </ul>
Phase III DUO-E NCT04269200	Advanced and recurrent endometrial cancer 1L	699	<ul style="list-style-type: none"> <li>Arm 1: chemo + <i>Imfinzi</i> placebo followed by <i>Imfinzi</i> placebo and <i>Lynparza</i> placebo</li> <li>Arm 2: chemo + <i>Imfinzi</i> followed by <i>Imfinzi</i> + <i>Lynparza</i> placebo</li> <li>Arm 3: chemo + <i>Imfinzi</i> followed by <i>Imfinzi</i> + <i>Lynparza</i></li> <li>Global Trial</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint PFS</li> <li>Secondary endpoints: OS, PFS2, ORR, DoR</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2020</li> <li>Data anticipated: 2023</li> </ul>
Phase II ORION NCT03775486	Stage IV NSCLC whose disease has not progressed following SoC chemo + <i>Imfinzi</i> maintenance therapy 1L	250	<ul style="list-style-type: none"> <li>Arm 1: <i>Imfinzi</i> + <i>Lynparza</i></li> <li>Arm 2: <i>Imfinzi</i> + placebo</li> <li>Global trial</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: PFS</li> <li>Secondary endpoints: OS, ORR, DoR, PFS in HRRm, PK, ADA</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2019</li> <li>Data readout: Q2 2021</li> </ul>
Phase I/II MEDIOLA NCT02734004	gBRCAm ovarian cancer 2L+ gBRCAm HER2-negative breast cancer 1-3L SCLC 2L+ Gastric cancer 2L+	148	<ul style="list-style-type: none"> <li>Dose until progression</li> <li>Arm 1: <i>Lynparza</i> + <i>Imfinzi</i></li> <li>Global trial</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: DCR at 12 weeks, safety and tolerability</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2016</li> <li>LPCD: Q2 2017</li> </ul>
Phase I/II MEDIOLA NCT02734004	gBRCAm ovarian cancer 2L+ Non-gBRCAm ovarian cancer 2L+ Non-gBRCAm ovarian cancer 2L+	115	<ul style="list-style-type: none"> <li>Dose until progression</li> <li>Arm 1: <i>Lynparza</i> + <i>Imfinzi</i></li> <li>Arm 2: <i>Lynparza</i> + <i>Imfinzi</i></li> <li>Arm 3: <i>Lynparza</i> + <i>Imfinzi</i> + bevacizumab</li> <li>Global trial</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: DCR at 12 weeks, ORR, safety and tolerability</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2018</li> <li>LPCD: Q2 2020</li> </ul>



# Lynparza (PARP inhibitor)

## Multiple cancers

Trial	Population	Patients	Design	Endpoints	Status
Phase III OlympiA NCT02032823 Partnered	BRCAm adjuvant breast cancer	1836	<ul style="list-style-type: none"> <li>Arm 1: <i>Lynparza</i> BID 12-month duration</li> <li>Arm 2: placebo 12-month duration</li> <li>Global trial partnership with Breast International Group and National Cancer Institute/NRG Oncology</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: invasive disease-free survival (iDFS)</li> <li>Secondary endpoint: distant disease-free survival and OS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2014</li> <li>LPCD: Q2 2019</li> <li>Data readout: Q1 2021</li> <li>Primary endpoint met</li> </ul>
Phase III MONO-OLA1 NCT04884360	BRCAwT advanced ovarian cancer 1L maintenance	420	<ul style="list-style-type: none"> <li>Arm 1: <i>Lynparza</i></li> <li>Arm 2: placebo</li> <li>Global trial, 12 countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: PFS (BRCAwT HRD+ve), PFS (BRCAwT)</li> <li>Secondary endpoints: OS, TFST, PFS2</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2021</li> <li>Data anticipated: &gt;2023</li> </ul>



# Lynparza (PARP inhibitor)

## Other combinations

Trial	Population	Patients	Design	Endpoints	Status
Phase III PROpel NCT03732820	Metastatic castration-resistant prostate cancer 1L	904	<ul style="list-style-type: none"> <li>Arm 1: <i>Lynparza</i> + abiraterone</li> <li>Arm 2: placebo + abiraterone</li> <li>Global trial, including China cohort</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: rPFS</li> <li>Secondary endpoints: OS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2018</li> <li>Data readout: Q3 2021</li> <li>Primary endpoint met</li> </ul>
Phase III LYNK-003 NCT04456699 Partnered	Advanced colorectal cancer 1L maintenance	525	<ul style="list-style-type: none"> <li>Arm 1: bevacizumab + 5-FU maintenance</li> <li>Arm 2: bevacizumab + <i>Lynparza</i> maintenance</li> <li>Arm 3: <i>Lynparza</i> maintenance</li> <li>Global trial</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: PFS</li> <li>Secondary endpoints: OS, ORR, DoR, AEs</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2020</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase II/III COCOS (GY005) NCT02502266 Partnered	Recurrent platinum resistant/refractory ovarian cancer	680	<ul style="list-style-type: none"> <li>Arm 1: chemo</li> <li>Arm 2: cediranib + <i>Lynparza</i></li> <li>Arm 3: cediranib</li> <li>Arm 4: <i>Lynparza</i></li> <li>US, Canada</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: PFS, OS</li> <li>Secondary endpoints: ORR, QoL, safety</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2016</li> <li>Data anticipated: 2023</li> </ul>
Phase II LYNK-002 NCT03742895 Partnered	HRRm or HRD-positive advanced cancer	390	<ul style="list-style-type: none"> <li>Arm 1: <i>Lynparza</i></li> <li>Global trial</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: ORR</li> <li>Secondary endpoints: DOR, OS, PFS, AE, Prog by CA-125</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2019</li> </ul>



# Enhertu (trastuzumab deruxtecan, HER2 ADC)

## Breast cancer

Trial	Population	Patients	Design	Endpoints	Status
Phase III DESTINY-Breast02 NCT03523585 Partnered	HER2-positive, unresectable and/or metastatic breast cancer pretreated with prior standard of care HER2 therapies, including trastuzumab emtansine	600	<ul style="list-style-type: none"> <li>Randomised open label parallel assignment</li> <li>Enhertu</li> <li>Physician's choice of lapatinib + capecitabine or trastuzumab + capecitabine</li> </ul>	<ul style="list-style-type: none"> <li>Primacy endpoint: PFS</li> <li>Secondary endpoints: OS, ORR, DoR, CBR</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2018</li> <li>LPCD: Q4 2020</li> <li>Data anticipated: H2 2022</li> </ul>
Phase III DESTINY-Breast03 NCT03529110 Partnered	HER2-positive, unresectable and/or metastatic breast cancer previously treated with trastuzumab and taxane	500	<ul style="list-style-type: none"> <li>Randomised open label parallel assignment</li> <li>Enhertu</li> <li>Ado-trastuzumab emtansine</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: PFS</li> <li>Secondary endpoints: OS, ORR, DoR, CBR</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2018</li> <li>LPCD: Q2 2020</li> <li>Data readout: Q3 2021</li> <li>Primary endpoint met</li> </ul>
Phase III DESTINY-Breast04 NCT03734029 Partnered	HER2-low, unresectable and/or metastatic breast cancer patients	540	<ul style="list-style-type: none"> <li>Randomised open label parallel assignment</li> <li>Enhertu</li> <li>Physician's choice of SoC chemo (choice of capecitabine, eribulin, gemcitabine, paclitaxel or nab-paclitaxel)</li> </ul>	<ul style="list-style-type: none"> <li>Primary end point: PFS</li> <li>Secondary endpoints: OS, DoR, ORR</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2018</li> <li>LPCD: Q4 2020</li> <li>Data readout: Q1 2022</li> <li>Primary endpoint met</li> </ul>
Phase III DESTINY-Breast05 NCT04622319 Partnered	High-risk HER2-positive patients with residual invasive breast cancer following neoadjuvant therapy	1600	<ul style="list-style-type: none"> <li>Randomised open label parallel assignment</li> <li>Enhertu</li> <li>Ado-trastuzumab emtansine</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: IDFS</li> <li>Secondary endpoints: DFS, OS, DRFI, BMFI</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2020</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase III DESTINY-Breast06 NCT04494425 Partnered	HER2-Low, HR+ breast cancer patients whose disease has progressed on endocrine therapy in the metastatic setting	850	<ul style="list-style-type: none"> <li>Randomised open label parallel assignment</li> <li>Enhertu</li> <li>Investigator's choice standard of care chemotherapy (capecitabine, paclitaxel, nab-paclitaxel)</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: PFS</li> <li>Secondary endpoints: OS, DoR, ORR</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2020</li> <li>Data anticipated: 2023</li> </ul>
Phase III DESTINY-Breast09 NCT04784715 Partnered	HER2-positive, metastatic breast cancer, no prior therapy for advanced or metastatic disease	1134	<ul style="list-style-type: none"> <li>Randomised, parallel assignment</li> <li>Enhertu + placebo</li> <li>Enhertu + pertuzumab</li> <li>Standard of care</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: PFS</li> <li>Secondary endpoints: OS, DoR, ORR</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2021</li> <li>Data anticipated: &gt;2023</li> </ul>



# Enhertu (trastuzumab deruxtecan, HER2 ADC)

## Breast cancer

Trial	Population	Patients	Design	Endpoints	Status
Phase III DESTINY-Breast11 NCT05113251 Partnered	High-risk HER2-positive early non-metastatic breast cancer	624	<ul style="list-style-type: none"> <li>Randomized open label parallel assignment</li> <li><i>Enhertu</i></li> <li><i>Enhertu</i>, followed by THP</li> <li>doxorubicin and cyclophosphamide, followed by THP</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: pCR</li> <li>Secondary endpoints: EFS, IDFS, OS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2021</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase Ib/II DESTINY-Breast07 NCT04538742 Partnered	HER2-positive metastatic breast cancer	450	<ul style="list-style-type: none"> <li>Randomised open label sequential assignment</li> <li><i>Enhertu</i></li> <li><i>Enhertu</i> + <i>Imfinzi</i></li> <li><i>Enhertu</i> + pertuzumab</li> <li><i>Enhertu</i> + paclitaxel</li> <li><i>Enhertu</i> + <i>Imfinzi</i> + paclitaxel</li> <li><i>Enhertu</i> + tucatinib</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: AE, SAE</li> <li>Secondary endpoints: ORR, PFS, DoR, OS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2021</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase Ib DESTINY-Breast08 NCT04556773 Partnered	HER2-low metastatic breast cancer	185	<ul style="list-style-type: none"> <li>Non-randomised open label parallel assignment</li> <li><i>Enhertu</i> + capecitabine</li> <li><i>Enhertu</i> + <i>Imfinzi</i> + paclitaxel</li> <li><i>Enhertu</i> + capivasertib</li> <li><i>Enhertu</i> + anastrozole</li> <li><i>Enhertu</i> + Faslodex</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: AE, SAE</li> <li>Secondary endpoints: ORR, PFS, DoR, OS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2021</li> <li>Data anticipated: 2023</li> </ul>



# Enhertu (trastuzumab deruxtecan, HER2 ADC)

## Gastric cancer

Trial	Population	Patients	Design	Endpoints	Status
Phase III DESTINY-Gastric04 NCT04704934 Partnered	HER2-positive gastric cancer or gastro-esophageal junction adenocarcinoma patients who have progressed on or after a trastuzumab-containing regimen and have not received any additional systemic therapy	490	<ul style="list-style-type: none"> <li>Open label randomised parallel group assignment</li> <li>Enhertu</li> <li>SoC chemo</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: OS</li> <li>Secondary endpoints: ORR, DoR, PFS, DcR, safety</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2021</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase II DESTINY-Gastric01 NCT03329690 Partnered	HER2-overexpressing advanced gastric or gastroesophageal junction adenocarcinoma patients who have progressed on two prior treatment regimens	233	<ul style="list-style-type: none"> <li>Randomised open label parallel assignment</li> <li>Enhertu</li> <li>SoC chemo</li> <li>Two additional open label patient cohorts with lower levels of HER2 expression</li> <li>Japan and Korea</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: ORR</li> <li>Secondary endpoints: PFS, OS, DoR, DCR, TTF, range of PK endpoints</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2017</li> <li>LPCD: Q2 2019</li> <li>Data readout: Q1 2020</li> <li>Primary endpoint met</li> </ul>
Phase II DESTINY-Gastric02 NCT04014075 Partnered	HER2-positive gastric cancer that cannot be surgically removed or has spread, in patients who have progressed on or after trastuzumab containing regimen	79	<ul style="list-style-type: none"> <li>Open label single group assignment</li> <li>Enhertu</li> <li>Western population</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: ORR</li> <li>Secondary endpoints: PFS, ORR, OS, DoR</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2019</li> <li>LPCD: Q4 2020</li> <li>Data readout: Q2 2021</li> <li>Primary endpoint met</li> </ul>
Phase II DESTINY-Gastric06 NCT04989816 Partnered	HER2-positive gastric cancer or gastro-esophageal junction adenocarcinoma patients who have progressed on two prior treatment regimens	75	<ul style="list-style-type: none"> <li>Open label single group assignment</li> <li>Enhertu</li> <li>China</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: ORR</li> <li>Secondary endpoints: PFS, ORR, DCR, OS, DoR, safety</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2021</li> <li>Data anticipated: 2023</li> </ul>
Phase Ib/II DESTINY-Gastric03 NCT04379596 Partnered	HER2-overexpressing gastric or gastroesophageal junction cancer patients	255	<ul style="list-style-type: none"> <li>Open label parallel assignment</li> <li>Part 1: To determine recommended Phase II combination dose</li> <li>5 Arms combine <i>Enhertu</i> with standard of care chemotherapies (5-FU, capecitabine, oxaliplatin) and / or durvalumab</li> <li>Part 2: To assess efficacy of the selected combinations</li> <li>Arm 2A: Standard chemotherapy (control)</li> <li>Arm 2B: <i>Enhertu</i> monotherapy</li> <li>Arm 2C: <i>Enhertu</i> with chemotherapy</li> <li>Arm 2D: <i>Enhertu</i> with chemotherapy and pembrolizumab</li> <li>Arm 2E: <i>Enhertu</i> and pembrolizumab</li> </ul>	<ul style="list-style-type: none"> <li>Part 1 Primary endpoint: safety</li> <li>Part 2 Primary endpoint: ORR</li> <li>Secondary endpoints: DoR, DCR, PFS, OS, range of PK endpoints, ADAs</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2020</li> <li>Data anticipated: 2023</li> </ul>



# Enhertu (trastuzumab deruxtecan, HER2 ADC)

## Other cancers

Trial	Population	Patients	Design	Endpoints	Status
Phase III DESTINY-Lung04 NCT05048797 Partnered	HER2 mutated, unresectable, locally advanced/metastatic NSCLC	264	<ul style="list-style-type: none"> <li>Randomised parallel group assignment</li> <li>Arm 1: <i>Enhertu</i></li> <li>Arm 2: SoC Treatment (platinum, pemetrexed and pembrolizumab)</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: PFS</li> <li>Secondary endpoints: OS, ORR, DoR, safety</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2021</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase II DESTINY-Lung01 NCT03505710 Partnered	HER2-over-expressing or mutated, unresectable and/or metastatic NSCLC	170	<ul style="list-style-type: none"> <li>Non randomised parallel group assignment</li> <li><i>Enhertu</i></li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: ORR</li> <li>Secondary endpoints: DoR, PFS, OS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2018</li> <li>Data readout: Q3 2021</li> <li>Primary endpoint met</li> </ul>
Phase II DESTINY-Lung02 NCT04644237 Partnered	HER2-mutated, unresectable and/or metastatic NSCLC	150	<ul style="list-style-type: none"> <li>Randomised parallel group assignment</li> <li>Arm 1: <i>Enhertu</i> 6.4 mg/kg</li> <li>Arm 2: <i>Enhertu</i> 5.4mg/kg</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: ORR</li> <li>Secondary endpoints: DoR, DCR, PFS, OS, PK</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2021</li> <li>Data anticipated: 2023</li> </ul>
Phase II DESTINY-PanTumour02 NCT04482309 Partnered	HER2 expressing tumours	280	<ul style="list-style-type: none"> <li>Non randomised single group assignment</li> <li><i>Enhertu</i></li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: ORR</li> <li>Secondary endpoints: DoR, DCR, PFS, OS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2020</li> <li>Data anticipated: 2023</li> </ul>
Phase II DESTINY-PanTumour01 NCT04639219 Partnered	HER2 mutant tumours	100	<ul style="list-style-type: none"> <li>Non-randomised single group assignment</li> <li><i>Enhertu</i></li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: ORR</li> <li>Secondary endpoints: DoR, DCR, PFS, PK</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2021</li> <li>Data anticipated: 2023</li> </ul>
Phase II DESTINY-CRC02 NCT04744831 Partnered	HER2 overexpressing advanced or metastatic colorectal cancer	120	<ul style="list-style-type: none"> <li>Randomised parallel group assignment</li> <li>Arm 1: <i>Enhertu</i> 6.4 mg/kg</li> <li>Arm 2: <i>Enhertu</i> 5.4mg/kg</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: ORR</li> <li>Secondary endpoint: PFS, OS, DoR, DCR, range of PK endpoints</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2021</li> <li>Data anticipated: 2023</li> </ul>



# *Enhertu* (trastuzumab deruxtecan, HER2 ADC)

## Other cancers

Trial	Population	Patients	Design	Endpoints	Status
Phase Ib DESTINY-Lung03 NCT04686305 Partnered	HER2-over-expressing, unresectable and/or metastatic NSCLC	136	<ul style="list-style-type: none"> <li>Non randomised parallel group assignment</li> <li>Part 1: To determine recommended combination dose</li> <li>3 Arms combine <i>Enhertu</i> with standard of care chemotherapies (cisplatin, carboplatin or pemetrexed) and <i>Imfinzi</i>. Arm 1D: <i>Enhertu</i> monotherapy arm</li> <li>Part 2: To assess efficacy of the selected combinations</li> <li>Arm 1: <i>Enhertu</i> + cisplatin + <i>Imfinzi</i></li> <li>Arm 2: <i>Enhertu</i> + carboplatin + <i>Imfinzi</i></li> <li>Arm 3: <i>Enhertu</i> + pemetrexed + <i>Imfinzi</i></li> <li>Arm 4: <i>Enhertu</i> + <i>Imfinzi</i></li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: safety</li> <li>Secondary endpoints: ORR, DoR, DCR, PFS, OS, range of PK endpoints</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2021</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase Ib U106 NCT04042701 Partnered	HER2 expressing locally advanced/metastatic breast or NSCLC	115	<ul style="list-style-type: none"> <li>Non randomised parallel group assignment</li> <li><i>Enhertu</i> + pembrolizumab</li> <li>Global trial 2 countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: DLT, ORR</li> <li>Secondary endpoints: DoR, DCR, PFS, TTR, OS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2020</li> </ul>
Phase Ib U105 NCT03523572 Partnered	HER2-expressing breast and urothelial cancer	99	<ul style="list-style-type: none"> <li>Non randomised sequential assignment</li> <li><i>Enhertu</i> + nivolumab</li> <li>Global trial 7 countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: DLT, ORR, TEAEs</li> <li>Secondary endpoints: DoR, DCR, PFS, TTR, OS, ORR (investigator)</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2018</li> <li>Data readout: Q3 2021</li> </ul>



# Calquence (BTK inhibitor)

## Blood and other cancers

Trial	Population	Patients	Design	Endpoints	Status
Phase I/II D8220C0007 NCT03932331	Chinese adults R/R MCL and R/R CLL	105	<ul style="list-style-type: none"> <li>Part 1: R/R B-cell Malignancies</li> <li>Part 2: Cohort A: R/R MCL</li> <li>Part 2: Cohort B: R/R CLL</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: safety, ORR</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2020</li> <li>Data anticipated: H2 2022</li> <li>First Data readout: MCL – Q4 2021, CLL – Q3 2022</li> <li>Second Data readout: CLL – Q1 2023</li> <li>Final data anticipated: MCL – Q3 2022, CLL – 2024+</li> </ul>
Phase I D8220C0001 NCT03198650	Japanese adults with advanced B-cell malignancies	35	<ul style="list-style-type: none"> <li>Part 1: Dose confirmation in r/r B-cell Malignancies</li> <li>Part 2: Dose expansion in r/r CLL or r/r MCL</li> <li>Calquence + obinutuzumab</li> <li>Part 3: Dose confirmation in treatment naïve CLL</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: safety</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2017</li> <li>Data anticipated: H2 2022</li> </ul>
Phase I D8220C00018 NCT04488016	Healthy volunteers	28	<ul style="list-style-type: none"> <li>Part 1: Rel bioavailability for capsule vs. tablet</li> <li>Part 2: Rel bioavailability for oral solution of tablet</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: safety, bioavailability</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2019</li> <li>Data readout: Q1 2021</li> <li>Primary endpoint met</li> </ul>
Phase I D8223C00013 NCT04768985	Healthy volunteers	66	<ul style="list-style-type: none"> <li>Arm A: Calquence tablet</li> <li>Arm B: Calquence capsule</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: bioequivalence</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2021</li> <li>Data readout: Q4 2021</li> <li>Primary endpoint met</li> </ul>



# Calquence (BTK inhibitor)

## Blood cancers

Trial	Population	Patients	Design	Endpoints	Status
Phase III ACE-CL-007 (ELEVATE-TN) NCT02475681	Previously untreated CLL	535	<ul style="list-style-type: none"> <li>Arm A: chlorambucil + obinutuzumab</li> <li>Arm B: <i>Calquence</i> + obinutuzumab</li> <li>Arm C: <i>Calquence</i></li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: PFS (Arm A vs. Arm B)</li> <li>Secondary endpoints: IRC (independent review committee) assessed ORR, OS (Arm A vs. Arm B vs. Arm C)</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2015</li> <li>Data readout: Q2 2019</li> <li>Primary endpoint met</li> </ul>
Phase III ACE-CL-311 NCT03836261	Previously untreated CLL fit	981	<ul style="list-style-type: none"> <li>Arm A: <i>Calquence</i> + venetoclax</li> <li>Arm B: <i>Calquence</i> + venetoclax + obinutuzumab</li> <li>Arm C: FCR or BR</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: IRC PFS (A vs C)</li> <li>Secondary endpoint: IRC PFS (B vs C); INV PFS (A vs C; B vs C)</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2019</li> <li>LPCD: Q4 2025</li> <li>Data anticipated: 2023</li> </ul>
Phase III ACE-CL-309 (ASCEND) NCT02970318	Relapsed/refractory CLL	306	<ul style="list-style-type: none"> <li>Arm A: <i>Calquence</i></li> <li>Arm B: rituximab + idelalisib or bendamustine (investigator's choice)</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: IRC assessed PFS (Arm A vs. Arm B)</li> <li>Secondary endpoints: INV-assessed ORR, OS, DoR, PROs</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2016</li> <li>Data readout: Q2 2019</li> <li>Primary endpoint met</li> </ul>
Phase III ACE-CL-006 (ELEVATE-RR) NCT02477696	Relapsed/refractory high risk CLL	533	<ul style="list-style-type: none"> <li>Arm A: <i>Calquence</i></li> <li>Arm B: ibrutinib</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: PFS</li> <li>Secondary endpoints: comparison of incidence of infections, RTs (Richter's Transformation) and atrial fibrillation, OS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2015</li> <li>Data readout: Q1 2021</li> <li>Primary endpoint met</li> </ul>
Phase III ACE-LY-308 (ECHO) NCT02972840	Previously untreated MCL	626	<ul style="list-style-type: none"> <li>Arm A: <i>Calquence</i> + bendamustine + rituximab</li> <li>Arm B: bendamustine + rituximab</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: PFS by Lugano Classification for NHL</li> <li>Secondary endpoints: IA, PFS, ORR, DoR, time to response, OS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2017</li> <li>LPCD: Q2 2023</li> <li>Data anticipated: 2023</li> </ul>
Phase III ESCALADE NCT04529772	DLBCL	600	<ul style="list-style-type: none"> <li><i>Calquence</i> + rituximab, cyclophosphamide, doxorubicin, vincristine, and prednisone</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: PFS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2021</li> <li>LPCD: Q3 2026</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase II 15-H-0016 NCT02337829	Relapsed/refractory and treatment naïve/del17p CLL/SLL	48	<ul style="list-style-type: none"> <li><i>Calquence</i> monotherapy</li> <li>Arm A: lymph node biopsy</li> <li>Arm B: bone marrow biopsy</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: ORR</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2014</li> <li>Data readout: Q4 2020</li> </ul>



# Calquence (BTK inhibitor)

## Blood cancers

Trial	Population	Patients	Design	Endpoints	Status
Phase Ib ACE-LY-106 NCT02717624	MCL	70	<ul style="list-style-type: none"> <li>• <i>Calquence</i> in combination with bendamustine and rituxumab</li> <li>• Arm A: treatment naïve</li> <li>• Arm B: relapsed/refractory</li> <li>• Arm C: treatment naïve: <i>Calquence</i> + venetoclax + rituxumab</li> </ul>	<ul style="list-style-type: none"> <li>• Primary endpoint: safety</li> </ul>	<ul style="list-style-type: none"> <li>• FPCD: Q1 2016</li> <li>• LPCD: Q2 2022</li> <li>• Data anticipated: H2 2022</li> </ul>
Phase I/II ACE-CL-001 NCT02029443	CLL/SLL/Richter's transformation	306	<ul style="list-style-type: none"> <li>• <i>Calquence</i> monotherapy</li> <li>• Dose escalation and expansion</li> </ul>	<ul style="list-style-type: none"> <li>• Primary endpoints: safety, PK, PD</li> </ul>	<ul style="list-style-type: none"> <li>• FPCD: Q1 2014</li> <li>• Data readout: Q4 2021</li> </ul>
Phase I ACE-LY-003 NCT02180711	Relapsed/refractory follicular lymphoma	80	<ul style="list-style-type: none"> <li>• Arm A: <i>Calquence</i></li> <li>• Arm B: <i>Calquence</i> + rituximab</li> <li>• Arm C: <i>Calquence</i> + rituximab + lenolidomide</li> </ul>	<ul style="list-style-type: none"> <li>• Primary endpoint: safety</li> </ul>	<ul style="list-style-type: none"> <li>• FPCD: Q1 2015</li> <li>• LPCD: Q3 2023</li> <li>• Data anticipated: &gt;2023</li> </ul>
Phase I ACE-CL-003 NCT02296918	CLL/SLL/PLL	69	<ul style="list-style-type: none"> <li>• <i>Calquence</i> + obinutuzumab</li> <li>• Arm A: relapsed/refractory</li> <li>• Arm B: treatment naïve</li> <li>• <i>Calquence</i> + venetoclax + rituxumab</li> <li>• Arm C: relapsed/refractory</li> <li>• Arm D: treatment naïve</li> </ul>	<ul style="list-style-type: none"> <li>• Primary endpoints: safety, ORR</li> <li>• Secondary endpoints: PD, PFS, TTNT, OS</li> </ul>	<ul style="list-style-type: none"> <li>• FPCD: Q4 2014</li> <li>• Data anticipated: H1 2022</li> </ul>



# Capivasertib (AKT inhibitor)

## Breast cancer, prostate cancer

Trial	Population	Patients	Design	Endpoints	Status
Phase III CAPitello-290 NCT03997123	Locally advanced or metastatic TNBC	924	<ul style="list-style-type: none"> <li>• Double-blind randomised comparative trial</li> <li>• Arm 1: capivasertib + paclitaxel</li> <li>• Arm 2: placebo + paclitaxel</li> </ul>	<ul style="list-style-type: none"> <li>• Primary endpoint: OS</li> </ul>	<ul style="list-style-type: none"> <li>• FPCD: Q3 2019</li> <li>• Data anticipated: 2023</li> </ul>
Phase III CAPitello-291 NCT04305496	2L and beyond in AI resistant locally advanced (Inoperable) or metastatic HR+/HER2- breast cancer	834	<ul style="list-style-type: none"> <li>• Double-blind randomised comparative trial</li> <li>• Arm 1: capivasertib + <i>Faslodex</i></li> <li>• Arm 2: placebo + <i>Faslodex</i></li> </ul>	<ul style="list-style-type: none"> <li>• Primary endpoint: PFS</li> </ul>	<ul style="list-style-type: none"> <li>• FPCD: Q2 2020</li> <li>• Data anticipated: H2 2022</li> </ul>
Phase III CAPitello-281 NCT04493853	De novo PTEN deficient metastatic hormone sensitive prostate cancer	1000	<ul style="list-style-type: none"> <li>• Double-blind randomised comparative trial</li> <li>• Arm 1: capivasertib + abiraterone</li> <li>• Arm 2: placebo + abiraterone</li> </ul>	<ul style="list-style-type: none"> <li>• Primary endpoint: rPFS</li> </ul>	<ul style="list-style-type: none"> <li>• FPCD: Q3 2020</li> <li>• Data anticipated: &gt;2023</li> </ul>
Phase III CAPitello-292 NCT04862663	1L triplet in early relapse/endocrine-resistant locally advanced (inoperable) or metastatic HR+/HER2- breast cancer	700	<ul style="list-style-type: none"> <li>• Double-blind randomised comparative trial</li> <li>• Arm 1: capivasertib + palbociclib + <i>Faslodex</i></li> <li>• Arm 2: placebo + palbociclib + <i>Faslodex</i></li> </ul>	<ul style="list-style-type: none"> <li>• Primary endpoint: PFS</li> </ul>	<ul style="list-style-type: none"> <li>• FPCD: Q2 2021</li> <li>• Data anticipated: &gt;2023</li> </ul>



# Camizestrant (AZD9833, next generation oral SERD)

## Breast cancer

Trial	Population	Patients	Design	Endpoints	Status
Phase III SERENA-4 NCT04711252	HR+ HER2- breast cancer	1342	<ul style="list-style-type: none"> <li>Randomised, double-blind, comparative trial</li> <li>Arm A: camizestrant (AZD9833) + palbociclib</li> <li>Arm B: anastrozole + palbociclib</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: PFS</li> <li>Secondary endpoint: OS, PFS2</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2021</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase III SERENA-6 NCT04964934	HR+ HER2- breast cancer	300	<ul style="list-style-type: none"> <li>Randomised, double-blind, comparator trial</li> <li>Arm A: camizestrant (AZD9833) plus palbociclib or abemaciclib</li> <li>Arm B: anastrozole or letrozole plus palbociclib or abemaciclib</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: PFS</li> <li>Secondary endpoint: OS, PFS2</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2021</li> <li>Data anticipated: 2023</li> </ul>
Phase II SERENA-2 NCT04214288	HR+ breast cancer	240	<ul style="list-style-type: none"> <li>Randomised, open-label, parallel-group, multicentre trial</li> <li>camizestrant (AZD9833) vs. i.m. Faslodex in women with advanced breast cancer.</li> </ul>	<ul style="list-style-type: none"> <li>Primary outcome: mPFS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2020</li> <li>LPCD: Q3 2021</li> <li>Data anticipated: H2 2022</li> </ul>
Phase II SERENA-3 NCT04588298	HR+ breast cancer	84	<ul style="list-style-type: none"> <li>Randomised, open-label, parallel-group, multicentre trial</li> </ul>	<ul style="list-style-type: none"> <li>Primary outcome: change in ER expression between pre- and on-treatment tumour biopsies</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2020</li> <li>Data anticipated: 2023</li> </ul>
Phase I NCT04541433	HR+ breast cancer	18	<ul style="list-style-type: none"> <li>Open-label</li> <li>anti-tumour activity of camizestrant (AZD9833) in Japanese women with endocrine resistant HR+ HER2- breast cancer that is not amenable to treatment with curative intent.</li> </ul>	<ul style="list-style-type: none"> <li>Primary outcomes: safety and tolerability</li> <li>Secondary outcome: PK</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2020</li> <li>LPCD: Q1 2022</li> <li>Data anticipated: 2023</li> </ul>
Phase I SERENA-1 NCT03616587	HR+ breast cancer	340	<ul style="list-style-type: none"> <li>Escalation phase - open label multicentre trial</li> <li>camizestrant (AZD9833)</li> <li>camizestrant (AZD9833) + palbociclib, everolimus, abemaciclib or capivasertib.</li> <li>Expansion phase - randomised expansion cohort(s)</li> <li>camizestrant (AZD9833)</li> <li>camizestrant (AZD9833) + palbociclib, everolimus, abemaciclib or capivasertib.</li> </ul>	<ul style="list-style-type: none"> <li>Primary outcomes: safety and tolerability</li> <li>Secondary outcomes: PK, antitumour activity</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2018</li> <li>Data anticipated: 2023</li> </ul>



# Camizestrant (AZD9833, next generation oral SERD)

## Breast cancer

Approved medicines

Late-stage development

Early development

Trial	Population	Patients	Design	Endpoints	Status
Phase I NCT04546347	Healthy volunteers	32	<ul style="list-style-type: none"><li>Randomised, open-label</li></ul>	<ul style="list-style-type: none"><li>Primary outcome: relative bioavailability of different tablet formulations and the effect of food</li></ul>	<ul style="list-style-type: none"><li>FPCD: Q3 2020</li><li>LPD: Q4 2020</li><li>Data readout: Q3 2021</li></ul>
Phase I NCT04818632	HR+ HER2- breast cancer Chinese patients	42	<ul style="list-style-type: none"><li>Dose escalation camizestrant (AZD9833)</li><li>Dose expansion camizestrant (AZD9833)</li><li>camizestrant (AZD9833) + palbociclib</li><li>camizestrant (AZD9833) + everolimus</li></ul>	<ul style="list-style-type: none"><li>Primary outcomes: safety and tolerability, PK</li><li>Secondary outcome: antitumour activity</li></ul>	<ul style="list-style-type: none"><li>FPCD: Q1 2021</li><li>Data anticipated: 2023</li></ul>

Oncology

CVRM

R&I

Other

V&I

Rare Disease



# Monalizumab (NKG2a mAb)

## Cancer

Approved medicines

Late-stage development

Early development

Oncology

CVRM

R&I

Other

V&I

Rare Disease

Trial	Population	Patients	Design	Endpoints	Status
Phase III INTERLINK-1 NCT04590963 Partnered	Recurrent or metastatic HNSCC, 2L	600	<ul style="list-style-type: none"><li>Arm A: monalizumab + cetuximab i.v.</li><li>Arm B: placebo + cetuximab i.v.</li><li>Global</li></ul>	<ul style="list-style-type: none"><li>Primary: OS</li><li>Secondary: PFS, ORR, DoR</li></ul>	<ul style="list-style-type: none"><li>FPCD: Q4 2020</li><li>Data anticipated: &gt;2023</li></ul>



# Datopotamab deruxtecan (TROP2 ADC)

## NSCLC

Trial	Population	Patients	Design	Endpoints	Status
Phase III TROPION-Lung01 NCT04656652 Partnered	Previously treated advanced or metastatic NSCLC with or without actionable genomic alterations	590	<ul style="list-style-type: none"> <li>Randomised, open label, parallel assignment</li> <li>Arm 1: datopotamab deruxtecan</li> <li>Arm 2: docetaxel</li> <li>Global</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: PFS, OS</li> <li>Secondary endpoints: ORR, DoR, TTR, DCR, PK, ADA</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2021</li> <li>Data anticipated: 2023</li> </ul>
Phase III TROPION-Lung08 NCT05215340 Partnered	Treatment-naïve patients with PD-L1 high advanced or metastatic NSCLC without actionable genomic alterations	740	<ul style="list-style-type: none"> <li>Randomised, open label</li> <li>Arm 1: datopotamab deruxtecan + pembrolizumab</li> <li>Arm 2: pembrolizumab</li> <li>Global</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: PFS, OS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2022</li> <li>Data anticipated: &gt;2023</li> <li>FSD achieved 8 March 2022</li> </ul>
Phase II TROPION-Lung05 NCT04484142 Partnered	Advanced or metastatic NSCLC with actionable genomic alterations and progressed on or after kinase inhibitor therapy and platinum-based chemotherapy	150	<ul style="list-style-type: none"> <li>Single-arm, open label</li> <li>datopotamab deruxtecan</li> <li>Global trial</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: ORR</li> <li>Secondary endpoints: DOR, PFS, OS, safety, PK, ADA</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2021</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase I TROPION-Lung02 NCT04526691 Partnered	Advanced or metastatic NSCLC	120	<ul style="list-style-type: none"> <li>Open label, two-part (dose escalation, dose expansion), sequential assignment</li> <li>datopotamab deruxtecan + pembrolizumab +/- platinum chemotherapy</li> <li>US, Japan</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: DLT, safety</li> <li>Secondary endpoints: ORR, DOR, PFS, OS, PK, ADA</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2020</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase I TROPION-Lung04 NCT04612751 Partnered	Advanced or metastatic NSCLC	120	<ul style="list-style-type: none"> <li>Open label, two-part (dose escalation, dose expansion), sequential assignment</li> <li>datopotamab deruxtecan + <i>Imfinzi</i> +/- platinum chemotherapy</li> <li>US, Japan</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: DLT, safety</li> <li>Secondary endpoints: ORR, DOR, PFS, OS, PK, ADA</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2021</li> <li>Data anticipated: &gt;2023</li> </ul>



# Datopotamab deruxtecan (TROP2 ADC)

## NSCLC and other cancers

Trial	Population	Patients	Design	Endpoints	Status
<b>Phase III</b> <b>TROPION-Breast01</b> <b>NCT05104866</b> <b>Partnered</b>	Inoperable or metastatic HR+, HER2-breast cancer	700	<ul style="list-style-type: none"> <li>Open-label, randomized</li> <li>datopotamab deruxtecan</li> <li>investigator's choice standard of care chemotherapy (eribulin, vinorelbine, capecitabine, gemcitabine)</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: PFS, OS</li> <li>Secondary endpoints: ORR, DoR, DCR, PL, ADA</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2021</li> <li>Data anticipated: &gt;2023</li> </ul>
<b>Phase I</b> <b>TROPION-PanTumor01</b> <b>NCT03401385</b> <b>Partnered</b>	Subjects with advanced solid tumours NSCLC TNBC HR+ BC HER2-negative gastric/GEJ esophageal Urothelial SCLC	770	<ul style="list-style-type: none"> <li>Open label, two-part (dose escalation, dose expansion), sequential assignment</li> <li>datopotamab deruxtecan</li> <li>Japan, US</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: DLT, safety</li> <li>Secondary endpoints: PK, anti-tumour activity, ADA</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2018</li> <li>Data anticipated: &gt;2023</li> <li>Early data readout (NSCLC) Q1 2021</li> <li>Early data readout (TNBC) Q2 2021</li> </ul>



# Calquence (BTK inhibitor)

## Blood cancers

Approved medicines

Late-stage development

Early development

Trial	Population	Patients	Design	Endpoints	Status
Phase III D822BC00001 NCT04075292	Untreated Chronic Lymphocytic Leukemia	150	<ul style="list-style-type: none"><li>Arm A: <i>Calquence</i></li><li>Arm B: Chlorambucil + rituximab</li></ul>	<ul style="list-style-type: none"><li>Primary endpoint: PFS</li><li>Secondary endpoint: ORR, DoR</li></ul>	<ul style="list-style-type: none"><li>FPCD: Q1 2015</li><li>LPCD: Q3 2022</li><li>Data anticipated: 2023</li><li>Final Data anticipated: Q4 2024</li></ul>

Oncology

CVRM

R&I

Other

V&I

Rare Disease




# Orpathys (savolitinib, MET inhibitor)

## NSCLC and other cancers

Trial	Population	Patients	Design	Endpoints	Status
Phase II NCT02897479 Partnered	Lung PSC and other NSCLC	65	<ul style="list-style-type: none"> <li>Single arm trial</li> <li>Orpathys QD</li> <li>China</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: ORR</li> <li>Secondary endpoints: PFS, safety parameters</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2017</li> <li>Data readout: Q2 2020</li> </ul>
Phase I NCT01985555 Partnered	Advanced NSCLC (all comers)	85	<ul style="list-style-type: none"> <li>Dose escalation trial</li> <li>China</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: safety and tolerability</li> <li>Secondary endpoints: PK profile</li> </ul>	<ul style="list-style-type: none"> <li>Data readout: Q3 2019</li> </ul>



# Oncology – early-stage development



# Adavosertib (WEE-1 inhibitor)

## Ovarian cancer, uterine serous cancer, solid tumours

Trial	Population	Patients	Design	Endpoints	Status
Phase II D6010C00004 NCT02272790	Platinum-resistant (PR) ovarian cancer	95	<ul style="list-style-type: none"> <li>Arm B: paclitaxel + adavosertib</li> <li>Arm C: carboplatin + adavosertib</li> <li>Global trial</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: ORR</li> <li>Secondary endpoints: DoR, PFS, OS, DCR, safety and tolerability</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2015</li> <li>LPCD: Q2 2018</li> <li>Data readout: Q3 2019</li> </ul>
Phase II D601HC00002 NCT04590248	Uterine serous carcinoma	120	<ul style="list-style-type: none"> <li>adavosertib monotherapy</li> <li>Phase IIb, open-label, single-arm, multicentre study</li> <li>Global trial</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: ORR</li> <li>Secondary endpoints: DoR, depth of response, PFS</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2020</li> <li>Data anticipated: H2 2022</li> </ul>
Phase I D6015C00002 NCT02617277	Advanced solid tumours	56	<ul style="list-style-type: none"> <li>Dose escalation trial</li> <li>adavosertib + <i>Imfinzi</i></li> <li>US</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: safety and tolerability, MTD</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2015</li> <li>LPCD: Q4 2018</li> <li>Data readout: Q4 2019</li> </ul>



# AZD0171 (anti-LIF mAb)

## Cancer

Approved medicines

Late-stage development

Early development

Oncology

Trial	Population	Patients	Design	Endpoints	Status
Phase II NCT04999969	1L metastatic pancreatic ductal adenocarcinoma	115	<ul style="list-style-type: none"><li>Open-label, non-randomised trial of AZD0171 + <i>Imfinzi</i> + gemcitabine, nab-paclitaxel</li></ul>	<ul style="list-style-type: none"><li>Primary endpoints: Safety, OS at 12m</li><li>Secondary endpoints: ORR, DoR, PFS</li></ul>	<ul style="list-style-type: none"><li>FPCD: Q1 2022</li><li>Data anticipated: &gt;2023</li></ul>

CVRM

R&I

Other

V&I

Rare Disease



# AZD0466 (Bcl2/xL inhibitor)

## Cancer

Approved medicines

Late-stage development

Early development

Oncology

CVRM

R&I

Other

V&I

Rare Disease

Trial	Population	Patients	Design	Endpoints	Status
Phase I/II NCT04865419	Advanced haematologic malignancies	160	<ul style="list-style-type: none"><li>Module 1</li><li>Part A: Dose escalation</li><li>AZD0466</li><li>Part B: Dose expansion</li><li>AZD0466</li><li>Module 2 - DDI trial</li><li>AZD0466 with voriconazole</li></ul>	<ul style="list-style-type: none"><li>Primary endpoint: safety</li><li>Secondary endpoint: PK</li></ul>	<ul style="list-style-type: none"><li>FPCD: Q2 2021</li><li>Data anticipated: &gt;2023</li></ul>



# AZD1390 (ATM inhibitor)

## Cancer

Approved medicines

Late-stage development

Early development

Oncology

Trial	Population	Patients	Design	Endpoints	Status
Phase I NCT03423628	Recurrent glioblastoma eligible for re-irradiation, brain metastases and leptomeningeal disease, newly-diagnosed glioblastoma patients	120	<ul style="list-style-type: none"><li>Open label trial</li><li>Arm A: recurrent GBM, AZD1390 +RT in dose escalation cohorts</li><li>Arm C: primary GBM, AZD1390 +RT in dose escalation cohorts</li></ul>	<ul style="list-style-type: none"><li>Primary endpoints: safety, tolerability, MTD</li><li>Secondary endpoints: PK and preliminary assessment of anti-tumour activity</li></ul>	<ul style="list-style-type: none"><li>FPCD: Q2 2018</li><li>Data anticipated: &gt;2023</li></ul>

CVRM

R&I

Other

V&I

Rare Disease



# AZD2936 (PD-1/TIGIT Bispecific mAb)

## Solid tumours

Trial	Population	Patients	Design	Endpoints	Status
<b>Phase I</b> <b>ARTEMIDE-01</b> <b>NCT04995523</b> <b>Partnered</b>	NSCLC	147	<ul style="list-style-type: none"> <li>Open-label, non-randomised dose-escalation and dose-expansion trial:</li> <li>Part A: Dose escalation in checkpoint inhibitor (CPI) experienced NSCLC pts with AZD2936 Intravenous monotherapy</li> <li>Part B: Dose expansion in checkpoint inhibitor (CPI) experienced NSCLC pts with AZD2936 intravenous monotherapy</li> <li>Part C: Dose expansion in CPI Naive NSCLC pts with AZD2936 i.v. monotherapy</li> <li>Part D: Dose expansion. Design to be confirmed by protocol amendment</li> <li>Europe, Australia, South Korea and North America</li> </ul>	<ul style="list-style-type: none"> <li>Part A Dose escalation primary endpoints: safety, RP2D, MTD</li> <li>Part B dose expansion primary endpoints: Safety and efficacy (ORR)</li> <li>Part C dose expansion primary endpoints: Safety and efficacy (ORR)</li> <li>Secondary endpoints: PK, PD (Receptor occupancy), efficacy (DCR, DoR, DRR, PFS)</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2021</li> <li>Data anticipated: &gt;2023</li> </ul>



# AZD4573 (CDK9 inhibitor)

## Blood cancers

Approved medicines

Late-stage development

Early development

Oncology

CVRM

R&I

Other

V&I

Rare Disease

Trial	Population	Patients	Design	Endpoints	Status
Phase II NCT05140382	R/R Peripheral T-cell lymphoma and R/R classical Hodgkin Lymphoma	90	<ul style="list-style-type: none"> <li>Open label, non-randomised modular dose confirmation and expansion study in pts with relapsed/ refractory PTCL or cHL</li> <li>Module 1: Exploring AZD4573 monotherapy treatment in 3 cohorts:                             <ul style="list-style-type: none"> <li>Cohort 1: PTCL, all comers (excluding NKTCL)</li> <li>Cohort 2: PTCL (NKTCL only)</li> <li>Cohort 3: cHL</li> </ul> </li> <li>i.v. Route of administration</li> <li>~30 centres in 10 countries across North America, Europe, EU and ROW</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: Efficacy</li> <li>Secondary endpoint: Safety, PK</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2022</li> <li>Data anticipated: 2023</li> </ul>
Phase I/II NCT04630756	R/R haematologic malignancies	90	<ul style="list-style-type: none"> <li>Open label, non-randomised trial</li> <li>Module 1 Part A: Dose setting</li> <li>AZD4573 + <i>Calquence</i> (100mg twice daily) combination in DLBCL, all comers; ramp-up across 3 dose levels</li> <li>Module 1 Part B: Dose expansion</li> <li>AZD4573 + <i>Calquence</i> (100mg twice daily) combination in GCB and non-GCB DLBCL</li> <li>Module 2 Part A: Dose Confirmation</li> <li>AZD4573 monotherapy window followed by AZD4573 + acalabrutinib in patients with relapsed or refractory MCL</li> <li>i.v. route of administration</li> <li>10 countries across North America, EU, ROW</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: Part A: safety</li> <li>Primary endpoint: Part B: ORR</li> <li>Secondary endpoints: Safety, PK, anti-tumour activity</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2021</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase I NCT03263637	R/R haematologic malignancies	44	<ul style="list-style-type: none"> <li>Arm 1: dose escalation in haematological malignancies excluding AML/ALL/high-risk MDS/CMML/CLL</li> <li>Arm 2: dose escalation in relapsed or refractory AML, ALL, high-risk MDS, CMML, CLL and Richter's syndrome</li> <li>i.v. route of administration</li> <li>Open in Netherlands, UK, Germany</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: safety, PK</li> <li>Secondary endpoint: efficacy</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2017</li> <li>LPD: Q3 2021</li> <li>Data anticipated: H2 2022</li> </ul>



# AZD5305 (PARP1 inhibitor)

## Solid tumours

Trial	Population	Patients	Design	Endpoints	Status
<b>Phase I/IIa</b> <b>PETRA</b> <b>NCT04644068</b>	Advanced, metastatic HER2- breast cancer (BRCAm, PALB2m or RAD51C/Dm) Advanced, metastatic TNBC PSR ovarian cancer (BRCAm, PALB2m or RAD51C/Dm) PSR ovarian cancer (HRD+) Prostate cancer (mCRPC, BRCAm) Prostate cancer (mCRPC, HRRm)	715	<ul style="list-style-type: none"> <li>• A modular, open-label, multicentre trial dose escalation</li> <li>• Module 1: AZD5305</li> <li>• Module 2: AZD5305 + paclitaxel</li> <li>• Module 3: AZD5305 + carboplatin +/- paclitaxel</li> <li>• Module 4: AZD5305 + trastuzumab deruxtecan</li> <li>• Module 5: AZD5305 + datopotamab deruxtecan</li> </ul>	<ul style="list-style-type: none"> <li>• Primary endpoints: safety and tolerability, PK</li> <li>• Secondary endpoints: efficacy</li> </ul>	<ul style="list-style-type: none"> <li>• FPCD: Q4 2020</li> <li>• Data anticipated: &gt;2023</li> </ul>



# AZD5991 (MCL1 inhibitor)

## Blood cancers

Approved medicines

Late-stage development

Early development

Trial	Population	Patients	Design	Endpoints	Status
Phase I/Ib/IIa NCT03218683	Relapsed/refractory haematologic malignancies	144	<ul style="list-style-type: none"><li>• Arm 1: AZD5991 dose escalation and expansion in R/R haematological malignancies</li><li>• Arm 2: AZD5991 + venetoclax combination dose escalation in R/R AML</li><li>• i.v. route of administration</li><li>• US only</li></ul>	<ul style="list-style-type: none"><li>• Primary endpoint: safety</li><li>• Secondary endpoints: PK, efficacy</li></ul>	<ul style="list-style-type: none"><li>• FPCD: Q3 2017</li><li>• Data anticipated: &gt;2023</li><li>• Suspended (The trial has been put on hold to allow further evaluation of safety related information)</li></ul>

Oncology

CVRM

R&I

Other

V&I

Rare Disease



# AZD7648 (selective DNA-PK inhibitor)

## Advanced solid tumours

Approved medicines

Late-stage development

Early development

Oncology

Trial	Population	Patients	Design	Endpoints	Status
Phase I NCT03907969	Advanced malignancies	192	<ul style="list-style-type: none"><li>Modular dose escalation and dose expansion trial</li><li>Arm 1: AZD7648 monotherapy</li><li>Arm 2: AZD7648 + pegylated liposomal doxorubicin</li><li>US, UK</li></ul>	<ul style="list-style-type: none"><li>Primary outcomes: safety and tolerability</li><li>Secondary outcomes: PK, Cytochrome P450, preliminary anti-tumour activity</li></ul>	<ul style="list-style-type: none"><li>FPCD: Q4 2019</li><li>Data anticipated: &gt;2023</li></ul>

CVRM

R&I

Other

V&I

Rare Disease



# AZD7789 (PD-1/TIM3 Bispecific mAb)

## Cancer

Trial	Population	Patients	Design	Endpoints	Status
Phase I/IIa NCT04931654	NSCLC Other tumours	81	<ul style="list-style-type: none"> <li>Open-label, non-randomised dose-escalation and dose-expansion trial:</li> <li>Part A: Dose escalation in post IO NSCLC pts with AZD7789 intravenous (iv) monotherapy</li> <li>Part B: Dose expansion in post IO and IO naïve NSCLC pts with AZD7789 iv monotherapy.</li> <li>North America, Europe</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: AE, SAE, DLTs, ORR</li> <li>Secondary endpoints: ORR, DCR, DoR, PFS, OS, PK, ADA</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2021</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase I/II NCT05216835	Relapsed or Refractory Classical Hodgkin Lymphoma	180	<ul style="list-style-type: none"> <li>Cohort A: Dose Escalation - Patients with anti-PD-1/PD-L1 exposed r/r cHL will receive AZD7789</li> <li>Cohort B1: Dose Expansion - Patients with anti-PD-1/PD-L1 exposed r/r cHL will receive AZD7789 once the recommended phase 2 dose (RP2D) has been determined</li> <li>Cohort B2: Dose Expansion - Patients with anti-PD-1/PD-L1 naïve r/r cHL will receive AZD7789 once the RP2D has been determined</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints:</li> <li>Cohort A: AE and DLTs</li> <li>Cohort B1: AE and ORR</li> <li>Cohort B2: AE and CRR</li> <li>Secondary endpoints:</li> <li>Cohort A: CRR, ORR, DoR, DoCR, PFS, OS, ADA and PK</li> <li>Cohort B1 and B2: DoR, DoCR, PFS, OS, ADA and PK</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2022</li> <li>Data anticipated: &gt;2023</li> </ul>



# AZD8205 (B7H4 ADC)

## Cancer

Approved medicines

Late-stage development

Early development

Oncology

Trial	Population	Patients	Design	Endpoints	Status
Phase I/II NCT05123482	Breast Cancer, Cholangiocarcinoma, Ovarian Cancer, Endometrial Cancer	198	<ul style="list-style-type: none"><li>Open-label, non-randomised dose-escalation and dose-expansion trial</li></ul>	<ul style="list-style-type: none"><li>Primary endpoints: AE, SAE, DLTs, Changes in Lab and PE parameters</li><li>Secondary endpoints: ORR, DCR, DoR, PFS, OS, PK, ADA</li></ul>	<ul style="list-style-type: none"><li>FPCD: Q1 2022</li><li>Data anticipated: &gt;2023</li></ul>

CVRM

R&I

Other

V&I

Rare Disease



# AZD8701 (FOXP3 antisense oligonucleotide)

## Solid tumours

Approved medicines

Late-stage development

Early development

Oncology

CVRM

R&I

Other

V&I

Rare Disease

Trial	Population	Patients	Design	Endpoints	Status
Phase I/Ib NCT04504669 Partnered	Advanced solid tumours	123	<ul style="list-style-type: none"><li>Dose escalation and dose expansion trial</li><li>Arm 1: AZD8701 monotherapy</li><li>Arm 2: AZD8701 &amp; Durvaluamb combination therapy</li><li>i.v. route of administration</li><li>US, CA, FR, ES</li></ul>	<ul style="list-style-type: none"><li>Primary endpoints: safety &amp; tolerability</li><li>Secondary endpoints: PK, PD, preliminary anti-tumour activity</li></ul>	<ul style="list-style-type: none"><li>FPCD: Q3 2020</li><li>Data anticipated: &gt;2023</li></ul>



# Ceralasertib (AZD6738, ATR inhibitor)

## Cancer

Approved medicines

Late-stage development

Early development

Oncology

CVRM

R&I

Other

V&I

Rare Disease

Trial	Population	Patients	Design	Endpoints	Status
Phase II PLANETTE NCT04564027	Solid tumours mCRPC	61	<ul style="list-style-type: none"> <li>Cohort A: ceralasertib; ATM-altered AST</li> <li>Cohort B: ceralasertib; ATM-altered mCRPC</li> <li>North America and Europe</li> </ul>	<ul style="list-style-type: none"> <li>Cohort A primary endpoint: ORR</li> <li>Cohort B primary endpoint: Composite RR</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2021</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase I NCT02264678	Solid tumours	330	<ul style="list-style-type: none"> <li>Module 1: ceralasertib + carboplatin</li> <li>Module 2: ceralasertib dose escalation, ceralasertib + <i>Lynparza</i></li> <li>Module 3: ceralasertib + <i>Imfinzi</i></li> <li>North America, Europe and South Korea</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: safety and tolerability.</li> <li>PK and efficacy</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2014</li> <li>Data anticipated: &gt;2023</li> </ul>



# IPH5201 (CD39 mAb)

## Solid tumours

Approved medicines

Late-stage development

Early development

Oncology

CVRM

R&I

Other

V&I

Rare Disease

Trial	Population	Patients	Design	Endpoints	Status
Phase I NCT04261075 Partnered	Advanced solid tumours	204	<ul style="list-style-type: none"><li>Open-label, dose-escalation trial to determine MTD of IPH5201 as monotherapy, or in combination with <i>Imfinzi</i> +/- oleclumab.</li><li>Part 1: IPH5201 monotherapy dose escalation to MTD</li><li>Part 2: IPH5201 + <i>Imfinzi</i> dose escalation to MTD</li><li>Part 3: IPH5201 + <i>Imfinzi</i> + oleclumab dose escalation to MTD</li><li>Route of administration: i.v.</li><li>4 countries - US and 3 in EU.</li></ul>	<ul style="list-style-type: none"><li>Primary endpoints: AE, SAE, DLT</li><li>Secondary endpoints: OR, DC, PK, ADA</li></ul>	<ul style="list-style-type: none"><li>FPCD: Q1 2020</li><li>Data anticipated: 2023</li></ul>



# MEDI1191 (IL-12 modRNA)

## Cancer

Approved medicines

Late-stage development

Early development

Oncology

CVRM

R&I

Other

V&I

Rare Disease

Trial	Population	Patients	Design	Endpoints	Status
Phase I NCT03946800 Partnered	Advanced solid tumours	87	<ul style="list-style-type: none"><li>Open-label, dose-escalation and expansion trial of MEDI1191 administered intratumourally as monotherapy and in combination with <i>Imfinzi</i></li></ul>	<ul style="list-style-type: none"><li>Primary endpoints: safety and tolerability</li><li>Secondary endpoints: PK, immunogenicity and efficacy</li></ul>	<ul style="list-style-type: none"><li>FPCD: Q2 2019</li><li>Data anticipated: &gt;2023</li></ul>



# MEDI5752 (PD-1/CTLA-4 bispecific mAb)

## Cancer

Trial	Population	Patients	Design	Endpoints	Status
Phase Ib NCT04522323	Advanced renal cell carcinoma	70	<ul style="list-style-type: none"> <li>Open-label, dose-escalation and dose-expansion trial</li> <li>Arm 1: MEDI5752 and axitinib</li> <li>Arm 2: MEDI5752 and lenvatinib</li> </ul>	<ul style="list-style-type: none"> <li>Dose escalation primary endpoints: safety, MTD, RP2D &amp; tolerability. Assess antitumour activity of the combination (ORR)</li> <li>Secondary endpoints: PK, ADA and antitumour activity (PFS, OR, DoR, DCR, TTR, OS)</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2020</li> <li>LPCD: Q2 2023</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase I NCT03530397	Advanced solid tumours	366	<ul style="list-style-type: none"> <li>Open-label, dose-escalation and dose-expansion trial</li> <li>Dose escalation: MEDI5752 i.v.</li> <li>Dose expansion: MEDI5752 i.v. as monotherapy and in combination with chemotherapy</li> <li>Arm A: MEDI5752 i.v.</li> <li>Arm B: MEDI5752 i.v., pemetrexed and carboplatin</li> <li>Arm C: pembrolizumab, pemetrexed and carboplatin</li> <li>Arm D: MEDI5752 i.v., taxane (paclitaxel or nab-paclitaxel) and carboplatin</li> </ul>	<ul style="list-style-type: none"> <li>Dose escalation primary endpoints: safety, tolerability, MTD, OBD and HPDD.</li> <li>Dose expansion primary endpoint: antitumour activity based on OR</li> <li>Secondary endpoints: PK, ADA, tumoral baseline PD-L1, antitumour activity (OR, DoR, DCR, PFS, OS)</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2018</li> <li>LPCD: Q3 2022</li> <li>Data anticipated: &gt;2023</li> </ul>



# MEDI9253 (rNDV-IL-12)

## Solid tumours

Approved medicines

Late-stage development

Early development

Oncology

CVRM

R&I

Other

V&I

Rare Disease

Trial	Population	Patients	Design	Endpoints	Status
Phase I NCT04613492	Advanced solid tumours	86	<ul style="list-style-type: none"><li>Open-label, dose-escalation and expansion trial</li><li>MEDI9253 + <i>Imfinzi</i></li></ul>	<ul style="list-style-type: none"><li>Primary endpoint: safety and tolerability</li><li>Secondary endpoints: PK, PD, immunogenicity and efficacy</li></ul>	<ul style="list-style-type: none"><li>FPCD: Q4 2020</li><li>Data anticipated: &gt;2023</li></ul>



# Oleclumab (CD73 mAb)

## Cancer

Approved medicines

Late-stage development

Early development

Oncology

CVRM

R&I

Other

V&I

Rare Disease

Trial	Population	Patients	Design	Endpoints	Status
Phase Ib/II Study 5 NCT03611556	Pancreatic 1L and 2L with prior gemcitabine-based chemotherapy	339	<ul style="list-style-type: none"> <li>Arm A1: gemcitabine and nab paclitaxel i.v.</li> <li>Arm A2: gemcitabine and nab paclitaxel i.v. + oleclumab i.v.</li> <li>Arm A3: gemcitabine and nab paclitaxel i.v. + oleclumab i.v. + <i>Imfinzi</i> i.v.</li> <li>Arm B1: mFOLFOX (oxaliplatin, leucovorin, 5-FU) i.v.</li> <li>Arm B2: mFOLFOX (oxaliplatin, leucovorin, 5-FU) i.v. + oleclumab i.v.</li> <li>Arm B3: mFOLFOX (oxaliplatin, leucovorin, 5-FU) i.v. + oleclumab i.v. + <i>Imfinzi</i> i.v.</li> <li>US, Norway, Spain and Australian trial centres</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: safety and anti-tumour activity</li> <li>Secondary endpoints include: PFS, PK, immunogenicity, safety and anti-tumour activity</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2018</li> <li>Data anticipated: 2023</li> </ul>



BioPharmaceuticals –  
approved medicines and  
late-stage pipeline



# Andexxa (anti-factor Xa reversal)

## Haematology

Approved medicines  
Late-stage development  
Early development

Trial	Population	Patients	Design	Endpoints	Status
Phase IV I8-513 (Post Launch) NCT03661528	Acute Intracranial Haemorrhage	1200	<ul style="list-style-type: none"> <li>Arm 1: <i>Andexxa</i></li> <li>Arm 2: Usual care</li> <li>Global trial</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: proportion of patients with good or excellent hemostatic efficacy as rated by an independent adjudication committee</li> <li>Secondary endpoint: change from baseline in anti-fXa activity</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2019</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase II 19-515 NCT04233073	Urgent surgery	10	<ul style="list-style-type: none"> <li>Arm 1: <i>Andexxa</i></li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: proportion of patients with good or excellent intraoperative hemostatic efficacy as determined by the surgeon's assessment and confirmed by an independent adjudication committee</li> <li>Secondary endpoint: percent change from baseline in anti-factor Xa activity</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2021</li> <li>LPCD: Q1 2022</li> <li>Data anticipated: H2 2022</li> </ul>

Oncology

CVRM

R&I

Other

V&I

Rare Disease



# Farxiga (SGLT2 inhibitor)

## Heart failure and chronic kidney disease

Trial	Population	Patients	Design	Endpoints	Status
<b>Phase III DELIVER NCT03619213</b>	CHF patients with HFpEF	6100	<ul style="list-style-type: none"> <li>Arm 1: <i>Farxiga</i> 10mg QD</li> <li>Arm 2: placebo</li> <li>Global trial - 21 countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: time to the first occurrence of any of the components of the composite: CV death or hospitalisation for HF or an urgent HF visit</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2018</li> <li>LPCD: Q1 2022</li> <li>Data anticipated: H1 2022</li> </ul>
<b>Phase III DAPA-MI NCT04564742</b>	Patients with myocardial infarction	6400	<ul style="list-style-type: none"> <li>Arm 1: <i>Farxiga</i> 10mg QD</li> <li>Arm 2: placebo</li> <li>Global trial - 2 countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: time to the first occurrence of any of the components of the composite: hospitalisation for HF or CV death</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2020</li> <li>Data anticipated: 2023</li> </ul>
<b>Phase I NCT04856007</b>	Healthy Chinese volunteers	80	<ul style="list-style-type: none"> <li>Arm 1: <i>Farxiga</i> 5 mg + metformin 500 mg XR</li> <li>Arm 2: <i>Farxiga</i>/metformin XR FDC 5/500 mg</li> <li>Arm 3: <i>Farxiga</i> 10 mg + metformin 1000 mg XR</li> <li>Arm 4: <i>Farxiga</i>/metformin XR FDC 10/1000 mg</li> <li>China only</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: Plasma AUCinf, AUClast and Cmax of <i>Farxiga</i> and metformin respectively.</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2021</li> <li>LPCD: Q2 2021</li> <li>Data readout: Q4 2021</li> </ul>



# Lokelma (sodium zirconium cyclosilicate)

## Hyperkalaemia

Trial	Population	Patients	Design	Endpoints	Status
<b>Phase IIIb DIALIZE China NCT04217590</b>	Patients with ESRD with hyperkalaemia and on stable haemodialysis	134	<ul style="list-style-type: none"> <li>Arm 1: <i>Lokelma</i> 5g QD for 8 weeks on non-dialysis days. Option to uptitrate to 10 and 15g QD.</li> <li>Arm 2: placebo QD for 8 weeks on non-dialysis days</li> <li>China</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: proportion of patients who maintain a pre-dialysis serum K between 4.0-5.0 mmol/L on 3 out of 4 dialysis treatments following the long interdialytic interval</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2020</li> <li>Data anticipated: H1 2022</li> </ul>
<b>Phase III HARMONIZE Asia NCT03528681</b>	Hyperkalaemia	250	<ul style="list-style-type: none"> <li>Open-label <i>Lokelma</i> 10g TID for 48 hours followed by:</li> <li>Arm 1: <i>Lokelma</i> 5g QD for 28 days</li> <li>Arm 2: <i>Lokelma</i> 10g QD for 28 days</li> <li>Arm 3: placebo QD for 28 days</li> <li>China</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: maintenance of normokalaemia</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2021</li> <li>Data anticipated: H2 2022</li> </ul>
<b>Phase III DIALIZE-Outcomes NCT04847232</b>	Patients with recurrent hyperkalaemia on chronic haemodialysis	2300	<ul style="list-style-type: none"> <li>Arm 1: <i>Lokelma</i> 5g-15g QD for 4 weeks on non-dialysis days, thereafter adjusted monthly</li> <li>Arm 2: placebo QD</li> <li>Global trial – 22 countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: Time to first occurrence of SCD, stroke, or hospitalisation/intervention/ED visit due to arrhythmias</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2021</li> <li>Data anticipated: &gt;2023</li> </ul>
<b>Phase III STABILIZE-CKD NCT05056727</b>	Patients with CKD and hyperkalaemia or at risk of hyperkalaemia	1360	<ul style="list-style-type: none"> <li>Open-label <i>Lokelma</i> (10g TID or 5g QD) for up to 72h, followed by 3 months open-label treatment with <i>Lokelma</i> (5g QOD to 15g QD) and uptitration of lisinopril or valsartan. Thereafter, patients are randomized to a 24 month treatment:</li> <li>Arm 1: <i>Lokelma</i> (5g QOD to 15g QD) and lisinopril or valsartan</li> <li>Arm 2: Placebo and lisinopril or valsartan</li> <li>Global trial – 13 countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: Total slope (eGFR measurements starting at randomisation) and chronic slope (eGFR measurements starting at 12 weeks after randomisation)</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2021</li> <li>Data anticipated: &gt;2023</li> </ul>



# Roxadustat (HIF-PH inhibitor)

## Anaemia

Trial	Population	Patients	Design	Endpoints	Status
<b>Phase III</b> <b>NCT03263091</b> <b>Partnered</b>	Anaemia in lower risk MDS patients	184	<ul style="list-style-type: none"> <li>Open label roxadustat lead-in</li> <li>Arm 1: roxadustat</li> <li>Arm 2: placebo</li> <li>US/global trial</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: proportion of patients achieving transfusion independence</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2017</li> <li>Data anticipated: 2023</li> <li>Sponsored by FibroGen</li> </ul>
<b>Phase II/III</b> <b>NCT03303066</b> <b>Partnered</b>	Anaemia in lower risk MDS patients	175	<ul style="list-style-type: none"> <li>Open label roxadustat lead-in</li> <li>Arm 1: roxadustat</li> <li>Arm 2: placebo</li> <li>China</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: haemoglobin response</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2018</li> <li>Data anticipated: 2023</li> <li>Sponsored by FibroGen</li> </ul>
<b>Phase II</b> <b>NCT04076943</b> <b>Partnered</b>	Anaemia in patients receiving chemotherapy treatment for non-myeloid malignancies	92	<ul style="list-style-type: none"> <li>Open label trial</li> <li>roxadustat 3x week</li> <li>16 weeks</li> <li>US</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: maximum change in haemoglobin within 16 weeks from baseline without RBC transfusion</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2019</li> <li>LPCD: Q3 2020</li> <li>Data readout: Q2 2021</li> <li>Primary endpoint met</li> <li>Sponsored by FibroGen</li> </ul>



# AZD4831 (MPO inhibitor)

## Cardiovascular disease

Trial	Population	Patients	Design	Endpoints	Status
Phase IIb/III NCT04986202	HFpEF	1485	<ul style="list-style-type: none"> <li>Randomised, double-blind</li> <li>Arm 1: 2.5mg AZD4831</li> <li>Arm 2: 5mg AZD4831</li> <li>Arm 3: placebo</li> <li>Global trial</li> </ul>	<ul style="list-style-type: none"> <li>Endpoints: Efficacy and Safety</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2021</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase IIa NCT03756285	HFpEF	96	<ul style="list-style-type: none"> <li>Arm 1: AZD4831</li> <li>Arm 2: placebo</li> <li>Global trial – five countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: The change from baseline in MPO activity in %</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2018</li> <li>Data readout: Q2 2021</li> </ul>
Phase I NCT02712372	Healthy patients	96	<ul style="list-style-type: none"> <li>SAD trial</li> <li>Arm 1: AZD4831</li> <li>Arm 2: placebo</li> <li>Germany</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: safety and tolerability</li> <li>Secondary endpoint: PK parameters</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2016</li> <li>LPCD: Q4 2016</li> <li>Data readout: Q2 2017</li> </ul>
Phase I NCT03136991	Healthy patients	40	<ul style="list-style-type: none"> <li>MAD trial</li> <li>Arm 1: AZD4831</li> <li>Arm 2: placebo</li> <li>USA</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: safety and tolerability</li> <li>Secondary endpoint: PK parameters</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2017</li> <li>LPCD: Q4 2017</li> <li>Data readout: Q1 2018</li> </ul>
Phase I NCT04232345	Healthy patients	32	<ul style="list-style-type: none"> <li>MAD trial in Japanese and Chinese patients</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: safety and tolerability</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2020</li> <li>Data readout: Q3 2021</li> </ul>
Phase I NCT04407091	Healthy patients	6	<ul style="list-style-type: none"> <li>Open label hADME trial</li> <li>a single oral dose of [<sup>14</sup>C] AZD4831</li> <li>UK</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: mass balance, with routes and rates of elimination of [<sup>14</sup>C]AZD4831; Metabolite profiling and structural identification; PK and total radioactivity</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2020</li> <li>LPCD: Q3 2020</li> <li>Data readout: Q3 2021</li> </ul>
Phase I NCT05052710	Healthy patients	14	<ul style="list-style-type: none"> <li>Open label</li> <li>AZD4831</li> <li>AZD4831 and Midazolam</li> <li>UK</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: PK parameters</li> <li>Secondary endpoints: safety and tolerability</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2021</li> <li>LPCD: Q4 2021</li> <li>Data anticipated: H1 2022</li> </ul>
Phase I NCT04949438	Renal Impairment	20	<ul style="list-style-type: none"> <li>Open label</li> <li>AZD4831</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: PK parameters</li> <li>Secondary endpoints: safety and tolerability</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2022</li> <li>Data anticipated: H2 2022</li> </ul>



# Eplontersen (ligand-conjugated antisense)

## ATTR

Approved medicines

Late-stage development

Early development

Oncology

CVRM

R&I

Other

V&I

Rare Disease

Trial	Population	Patients	Design	Endpoints	Status
<b>Phase III CARDIO-TTRansform NCT04136171</b> Partnered with Ionis Pharmaceuticals Inc	Patients with hereditary or wild-type Transthyretin-Mediated Amyloid Cardiomyopathy (ATTR CM)	750	<ul style="list-style-type: none"> <li>Arm 1: eplontersen s.c.</li> <li>Arm 2: placebo</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: composite Outcome of Cardiovascular (CV) Mortality and recurrent CV clinical events at Week 120</li> <li>Secondary endpoints include: 6MWT, KCCQ, CV Events, CV Mortality</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2020</li> <li>Data anticipated: &gt;2023</li> </ul>
<b>Phase III NEURO-TTRansform NCT04136184</b> Partnered with Ionis Pharmaceuticals Inc	Patients with hereditary transthyretin-mediated amyloid polyneuropathy (hATTR-PN)	168	<ul style="list-style-type: none"> <li>Arm 1: eplontersen s.c.</li> <li>Arm 2: inotersen s.c.</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: change from baseline in mNIS+7 at Week 66</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2020</li> <li>Data anticipated: H1 2022</li> </ul>



# Breztri (PT010, LAMA/LABA/ICS, pMDI)

## Asthma

Trial	Population	Patients	Design	Endpoints	Status
Phase III KALOS NCT04609878	Severe asthma	2200	<ul style="list-style-type: none"> <li>Randomised, double-blind, double dummy, parallel group and multicentre</li> <li>Treatments (24 to 52 week variable length)</li> <li>BGF 320/28.8/9.6µg BID MDI</li> <li>BGF 320/14.4/9.6µg BID MDI</li> <li>BFF 320/9.6µg BID MDI</li> <li>Symbicort 320/9µg BID pMDI</li> <li>Multi-country</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: Change from baseline in forced expiratory volume in 1 second (FEV1) area under the curve 0 to 3 hours (AUC0-3) at Week 24</li> <li>Primary endpoint of pooled trials D5982C00007 and D5982C00008: Rate of severe asthma exacerbations</li> <li>Secondary endpoint: Change from baseline in morning pre-dose trough FEV1 at Week 24</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2021</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase III LOGOS NCT04609904	Severe asthma	2200	<ul style="list-style-type: none"> <li>Randomised, double-blind, double dummy, parallel group and multicentre</li> <li>Treatments (24 to 52 week variable length)</li> <li>BGF 320/28.8/9.6µg BID MDI</li> <li>BGF 320/14.4/9.6µg BID MDI</li> <li>BFF 320/9.6µg BID MDI</li> <li>Symbicort 320/9µg BID pMDI</li> <li>Multi-country</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: Change from baseline in forced expiratory volume in 1 second (FEV1) area under the curve 0 to 3 hours (AUC0-3) at Week 24</li> <li>Primary endpoint of pooled trials D5982C00007 and D5982C00008: Rate of severe asthma exacerbations</li> <li>Secondary endpoint: Change from baseline in morning pre-dose trough FEV1 at Week 24</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2021</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase III VATHOS NCT05202262	Moderate asthma	630	<ul style="list-style-type: none"> <li>Randomized, double-blind, parallel group and multicentre</li> <li>Treatments (24 week)</li> <li>BFF 320/9.6µg BID MDI</li> <li>BFF 160/9.6µg BID MDI</li> <li>BD 320µg BID MDI</li> <li>Open-label Symbicort TBH 320/9µg BID</li> <li>Multi-country</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: Change from baseline in forced expiratory volume in 1 second (FEV1) area under the curve 0 to 3 hours (AUC0-3) at Week 24</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2022</li> <li>Data anticipated: &gt;2023</li> </ul>



# *Daliresp/ Daxas* (PDE4 inhibitor, oral)

## COPD

Trial	Population	Patients	Design	Endpoints	Status
Phase IV PASS (post launch) NCT03381573	COPD	124080	<ul style="list-style-type: none"> <li>A retrospective cohort trial comparing COPD patients aged 40 years and older with new exposure to roflumilast with up to 5 unexposed (i.e., not roflumilast-exposed) COPD controls matched by propensity score (PS), age, sex, and year of cohort entry.</li> <li>The trial is using electronic healthcare databases in the US (Military Health System database), Germany (German Pharmacoepidemiological Research Database), Sweden (national databases including healthcare, death, and demographics data) and Norway.</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: all-cause mortality (up to five years)</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2017</li> <li>LPCD: Q3 2022</li> <li>Data anticipated: H2 2022</li> </ul>



# Fasenra (IL-5R mAb)

## Dermatology

Approved medicines  
Late-stage development  
Early development

Trial	Population	Patients	Design	Endpoints	Status
<b>Phase III FJORD NCT04612790</b>	Patients with symptomatic (newly diagnosed or relapsing) bullous pemphigoid	120	<ul style="list-style-type: none"> <li>• Double blind treatment period and open label period</li> <li>• Arm 1: <i>Fasenra</i></li> <li>• Arm 2: placebo</li> <li>• 36-week</li> <li>• Global trial</li> </ul>	<ul style="list-style-type: none"> <li>• Primary endpoint: Proportion of patients with complete sustained (<math>\geq 2</math> months) remission off OCS at 36 weeks</li> </ul>	<ul style="list-style-type: none"> <li>• FPCD: Q2 2021</li> <li>• Data anticipated: 2023</li> </ul>
<b>Phase II ARROYO NCT04612725</b>	Patients with moderate/severe chronic spontaneous urticaria, and resistant to H1 treatment	160	<ul style="list-style-type: none"> <li>• Double blind treatment period and open label period</li> <li>• Arm 1: <i>Fasenra</i> regimen 1</li> <li>• Arm 2: <i>Fasenra</i> regimen 2</li> <li>• Arm 3: placebo</li> <li>• 24-week</li> <li>• Global trial</li> </ul>	<ul style="list-style-type: none"> <li>• Primary endpoint: Change from baseline in ISS7 at week 12</li> </ul>	<ul style="list-style-type: none"> <li>• FPCD: Q4 2020</li> <li>• Data anticipated: H2 2022</li> </ul>
<b>Phase II HILLIER NCT04605094</b>	Patients with moderate to severe atopic dermatitis despite treatment with topical medications	194	<ul style="list-style-type: none"> <li>• Double blind treatment period and open label periods</li> <li>• Arm 1: <i>Fasenra</i></li> <li>• Arm 2: placebo</li> <li>• 16-week DB period</li> <li>• Global trial</li> </ul>	<ul style="list-style-type: none"> <li>• Primary endpoint: Proportion of patients with an IGA 0/1 and a decrease in IGA of <math>\geq 2</math> points at week 16</li> </ul>	<ul style="list-style-type: none"> <li>• FPCD: Q4 2020</li> <li>• LPCD: Q4 2021</li> <li>• Data anticipated: H2 2022</li> </ul>

Oncology

CVRM

R&I

Other

V&I

Rare Disease



# Fasenra (IL-5R mAb)

## Gastrointestinal diseases

Approved medicines

Late-stage development

Early development

Trial	Population	Patients	Design	Endpoints	Status
<b>Phase III MESSINA NCT04543409</b>	Documented diagnosis of EoE Age 12 to 65 years	170	<ul style="list-style-type: none"> <li>• Double blind treatment period and open label period(s)</li> <li>• Arm 1: <i>Fasenra</i> 30mg Q4W s.c.</li> <li>• Arm 2: placebo Q4W s.c.</li> <li>• 24-week</li> <li>• Global trial – 12 countries</li> </ul>	<ul style="list-style-type: none"> <li>• Primary endpoints: histologic response at week 24, change from baseline in DSQ score at week 24</li> </ul>	<ul style="list-style-type: none"> <li>• FPCD: Q4 2020</li> <li>• Data anticipated: H2 2022</li> </ul>
<b>Phase III HUDSON NCT05251909</b>	Patients with eosinophilic gastritis and/or gastroenteritis. Age >=12yrs	220	<ul style="list-style-type: none"> <li>• Double blind treatment period and open label extension</li> <li>• Arm 1: <i>Fasenra</i> s.c.</li> <li>• Arm 2: placebo s.c.</li> <li>• 24-week</li> <li>• Global trial</li> </ul>	<ul style="list-style-type: none"> <li>• Dual primary endpoints at week 24:</li> <li>• Proportion of patients achieving a histological response in the stomach and/or in the duodenum</li> <li>• Absolute change in symptoms of EG/EGE</li> </ul>	<ul style="list-style-type: none"> <li>• FPCD: Q1 2022</li> <li>• Data anticipated: &gt;2023</li> </ul>

Oncology

CVRM

R&I

Other

V&I

Rare Disease



# Fasenra (IL-5R mAb)

## Nasal polyposis and other eosinophilic diseases

Trial	Population	Patients	Design	Endpoints	Status
<b>Phase III OSTRO NCT03401229</b>	Patients with severe bilateral nasal polyps who are still symptomatic despite standard of care therapy Age 18-75 years	413	<ul style="list-style-type: none"> <li>Arm 1: <i>Fasenra</i> 30mg Q8W s.c.</li> <li>Arm 2: placebo s.c.</li> <li>56-week trial</li> <li>Global trial- 8 countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: effect of <i>Fasenra</i> on nasal polyp burden and on patient reported nasal blockage</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2018</li> <li>LPCD: Q2 2019</li> <li>Data readout: Q3 2020</li> <li>Co-primary endpoints met</li> </ul>
<b>Phase III ORCHID NCT04157335</b>	Patients with eosinophilic chronic rhinosinusitis with severe nasal polyposis Age 18-75 years	148	<ul style="list-style-type: none"> <li>Arm 1: <i>Fasenra</i> 30mg Q8W s.c.</li> <li>Arm 2: placebo Q8W s.c.</li> <li>56-week trial</li> <li>Global trial - 10 countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: Change in endoscopic total nasal polyp score and change in mean nasal blockage score</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2019</li> <li>Data anticipated: 2023</li> </ul>
<b>Phase III MANDARA NCT04157348</b>	Patients with relapsing or refractory EGPA on corticosteroid therapy with or without stable immunosuppressive therapy Age 18 years and older	140	<ul style="list-style-type: none"> <li>Arm 1: <i>Fasenra</i> 30mg Q4W s.c.</li> <li>Arm 2: mepolizumab 300mg Q4W s.c.</li> <li>52-week trial with a minimum 1-year open label extension</li> <li>Global trial- 9 countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: Proportion of patients achieving remission (BVAS=0 and OCS dose <math>\leq</math> 4mg/day) at both weeks 36 and 48.</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2019</li> <li>Data anticipated: 2023</li> </ul>
<b>Phase III NATRON NCT04191304</b>	Patients with HES (history of persistent eosinophilia >1500 cells/ $\mu$ L with evidence of end organ manifestations attributable to eosinophilia) and signs or symptoms of HES worsening/flare at visit 1 Age 12 years and older	120	<ul style="list-style-type: none"> <li>Arm 1: <i>Fasenra</i> 30mg Q4W s.c.</li> <li>Arm 2: placebo Q4W s.c.</li> <li>24-week trial with a minimum 1-year open label extension</li> <li>Global trial- 9-12 countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: Time to first HES worsening/flare</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2020</li> <li>Data anticipated: 2023</li> </ul>



# Fasenra (IL-5R mAb)

## Severe, uncontrolled asthma

Trial	Population	Patients	Design	Endpoints	Status
Phase IIIb PONENTE NCT03557307	Severe eosinophilic asthmatics receiving HD ICS + LABA and chronic OCS with or without additional asthma controller(s). Age 18 Years and older	599	<ul style="list-style-type: none"> <li>Arm 1: <i>Fasenra</i> 30mg Q8W s.c.</li> <li>38-week trial</li> <li>Global trial – 16 countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: reduction of oral corticosteroid dose</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2018</li> <li>LPCD: Q3 2019</li> <li>Data readout: Q4 2020</li> <li>Primary endpoint met</li> </ul>
Phase III D3250C00036 China ICS/LABA Trial (MIRACLE) NCT03186209	Severe, uncontrolled asthma, despite background controller medication, MD & HD ICS + LABA ± chronic OCS Age 12-75 years	695	<ul style="list-style-type: none"> <li>Arm 1: <i>Fasenra</i> 30mg Q8W s.c.</li> <li>Arm 2: placebo s.c.</li> <li>56-week trial</li> <li>Global trial – 4 countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: annual asthma exacerbation rate</li> <li>Secondary endpoints: assess pulmonary function, asthma symptoms, other asthma control metrics</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2017</li> <li>LPCD: Q4 2021</li> <li>Data anticipated: 2023</li> </ul>



# Fasenra (IL-5R mAb)

## Severe, uncontrolled asthma, COPD and other eosinophilic diseases

Trial	Population	Patients	Design	Endpoints	Status
<b>Phase III RESOLUTE NCT04053634</b>	Patients with moderate to very severe COPD with a history of frequent exacerbations on a background triple therapy (ICS/LABA/LAMA) Age 40-85 years	868	<ul style="list-style-type: none"> <li>• Double-blind, placebo controlled</li> <li>• Arm 1: <i>Fasenra</i> 100mg Q8W s.c.</li> <li>• Arm 2: placebo Q8W s.c.</li> <li>• 56-week treatment</li> <li>• Global trial – 26 countries</li> </ul>	<ul style="list-style-type: none"> <li>• Primary endpoint: annualized rate of moderate or severe exacerbations over 56 weeks</li> </ul>	<ul style="list-style-type: none"> <li>• FPCD: Q4 2019</li> <li>• Data anticipated: &gt;2023</li> </ul>
<b>Phase III MAHALE NCT05006573</b>	Patients With non-cystic fibrosis bronchiectasis (NCFB) with eosinophilic inflammation Age 18 years and older	420	<ul style="list-style-type: none"> <li>• Double blind treatment period and open label extension trial</li> <li>• Arm 1: <i>Fasenra</i> 30mg Q4W s.c.</li> <li>• Arm 2: placebo Q4W s.c.</li> <li>• 52-week</li> <li>• Global trial – 17 countries</li> </ul>	<ul style="list-style-type: none"> <li>• Primary endpoint: annualised bronchiectasis exacerbation rate at week 52</li> </ul>	<ul style="list-style-type: none"> <li>• FPCD: Q3 2021</li> <li>• Data anticipated: &gt;2023</li> </ul>



# Saphnelo (type I interferon receptor mAb)

## Lupus (SLE / LN)

Approved medicines  
Late-stage development  
Early development

Trial	Population	Patients	Design	Endpoints	Status
Phase III TULIP LTE NCT02794285	Moderate to severe SLE	630	<ul style="list-style-type: none"> <li>Arm 1: 300mg i.v. anifrolumab Q4W for 152 weeks</li> <li>Arm 2: placebo i.v. Q4W for 152 weeks</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: extension to evaluate long-term safety and tolerability</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2016</li> <li>LPCD: Q4 2018</li> <li>Data anticipated: H1 2022</li> </ul>
Phase III TULIP-SC NCT04877691 Partnered	Moderate to severe SLE patients	360	<ul style="list-style-type: none"> <li>Arm 1: <i>Saphnelo</i> s.c.</li> <li>Arm 2: placebo s.c.</li> <li>Global</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: BICLA at week 52</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2021</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase III AZALEA-SLE NCT04931563 Partnered	Moderate to severe SLE patients	328	<ul style="list-style-type: none"> <li>Arm 1: 300 mg <i>Saphnelo</i> i.v. Q4W</li> <li>Arm 2: 300 mg placebo i.v. Q4W</li> <li>Asia</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: BICLA at week 52</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2021</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase II TULIP-LN1 NCT02547922	Active proliferative LN	150	<ul style="list-style-type: none"> <li>Arm 1: 900 mg i.v. Q4W for 12 weeks then 300mg i.v. anifrolumab Q4W for 36 weeks</li> <li>Arm 2: 300 mg i.v. anifrolumab Q4W for 48 weeks</li> <li>Arm 3: placebo i.v. Q4W for 48 weeks</li> </ul>	<ul style="list-style-type: none"> <li>Response in proteinuria at week 52</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2015</li> <li>LPCD: Q4 2018</li> <li>Primary endpoint not met</li> </ul>

Oncology

CVRM

R&I

Other

V&I

Rare Disease



# Tezspire (TSLP mAb)

## Severe, uncontrolled asthma

Trial	Population	Patients	Design	Endpoints	Status
<b>Phase III NAVIGATOR</b> NCT03347279 Partnered	Severe asthma Age 12-80 years	1061	<ul style="list-style-type: none"> <li>Arm 1: Tezspire s.c.</li> <li>Arm 2: placebo s.c.</li> <li>52 week trial</li> <li>Global trial – 18 countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: Annual asthma exacerbation rate</li> <li>Secondary endpoints: Change from baseline in pre-BD FEV1, asthma related QoL (AQLQ(S)+12), asthma control (ACQ-6)</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2018</li> <li>LPCD: Q3 2019</li> <li>Data readout: Q4 2020</li> <li>Primary endpoint met</li> </ul>
<b>Phase III SOURCE</b> NCT03406078 Partnered	Severe asthma Age 18-80 years	150	<ul style="list-style-type: none"> <li>Arm 1: Tezspire s.c.</li> <li>Arm 2: placebo s.c.</li> <li>48 week trial</li> <li>Global trial – seven countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: Reduction from baseline in daily OCS dose while not losing asthma control</li> <li>Secondary endpoint: Annual asthma exacerbation rate</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2018</li> <li>LPCD: Q4 2019</li> <li>Data readout: Q4 2020</li> <li>Primary endpoint not met</li> </ul>
<b>Phase III DESTINATION</b> NCT03706079 Partnered	Severe asthma Age 12-80 years	951	<ul style="list-style-type: none"> <li>Extension trial to NAVIGATOR and SOURCE</li> <li>Arm 1: Tezspire s.c.</li> <li>Arm 2: placebo s.c.</li> <li>52 week trial (subjects from NAVIGATOR); 56 week trial (subjects from SOURCE)</li> <li>Global trial – 18 countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: Exposure adjusted rates of AEs/SAEs</li> <li>Secondary endpoints: Annual asthma exacerbation rate</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2019</li> <li>LPCD: Q4 2020</li> <li>Data anticipated: H2 2022</li> </ul>
<b>Phase III PATH-HOME</b> NCT03968978 Partnered	Severe asthma Age 12-80 years	216	<ul style="list-style-type: none"> <li>Arm 1: Tezspire s.c. via AI</li> <li>Arm 2: Tezspire s.c. via APFS</li> <li>24 week trial</li> <li>Global trial – 4 countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: Proportion of health care professionals and patients /caregivers who successfully administered Tezspire in clinic and at home with an APFS or an AI, respectively</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2019</li> <li>LPCD: Q3 2019</li> <li>Data readout: Q4 2020</li> <li>Primary endpoint met</li> </ul>



# Tezspire (TSLP mAb)

## Severe, uncontrolled asthma, COPD & CRSwNP

Trial	Population	Patients	Design	Endpoints	Status
Phase III WAYPOINT NCT04851964 Partnered	Severe chronic rhinosinusitis with nasal polyps (CRSwNP) Age 18+	400	<ul style="list-style-type: none"> <li>Arm 1: Tezspire s.c.</li> <li>Arm 2: placebo s.c.</li> <li>52 week trial</li> <li>Global trial – 10 countries</li> </ul>	<ul style="list-style-type: none"> <li>Co-primary endpoint: nasal polyp score and participant reported nasal congestion</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2021</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase III DIRECTION NCT03927157 Partnered	Severe asthma Age 18-80 years	396	<ul style="list-style-type: none"> <li>Arm 1: Tezspire s.c.</li> <li>Arm 2: placebo s.c.</li> <li>52 week trial</li> <li>Regional Asia trial – three countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: annual asthma exacerbation rate</li> <li>Secondary endpoints: change from baseline in pre-BD FEV1, asthma related QoL (AQLQ(S)+12), asthma control (ACQ-6)</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2019</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase III NOZOMI NCT04048343 Partnered	Severe asthma 12-80 years	65	<ul style="list-style-type: none"> <li>Arm 1: Tezspire s.c.</li> <li>52 week trial</li> <li>Local trial - Japan</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: number of patients with adverse events</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2019</li> <li>LPCD: Q1 2020</li> <li>Data readout: Q2 2021</li> <li>Primary endpoint met</li> </ul>
Phase IIa COURSE NCT04039113 Partnered	Moderate to very severe COPD Age 40-80	338	<ul style="list-style-type: none"> <li>Arm 1: Tezspire s.c.</li> <li>Arm 2: placebo s.c.</li> <li>52 week trial</li> <li>Global trial – 10 countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: rate of moderate or severe COPD exacerbations</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2019</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase II CASCADE NCT03688074 Partnered	Severe asthma Age 18-75 years	116	<ul style="list-style-type: none"> <li>Arm 1: Tezspire s.c.</li> <li>Arm 2: placebo s.c.</li> <li>28 week trial</li> <li>Global trial – five countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: number of airway submucosal inflammatory cells/mm<sup>2</sup> of bronchoscopic biopsies</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2018</li> <li>LPCD: Q4 2019</li> <li>Data readout: Q2 2021</li> <li>Primary endpoint met</li> </ul>



# Brazikumab (IL-23 inhibitor)

## Inflammatory bowel disease (Crohn's disease, ulcerative colitis)

Trial	Population	Patients	Design	Endpoints	Status
<b>Phase III NCT03961815</b>	Crohn's Disease	161	<ul style="list-style-type: none"> <li>An Open-label, Long-term Extension Safety Study of Brazikumab in Participants With Moderately to Severely Active Crohn's Disease</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: safety of long-term treatment with Brazikumab (AEs, clinical laboratory values, vital signs, and ECGs)</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2020</li> <li>Data anticipated: &gt;2023</li> </ul>
<b>Phase IIb/III INTREPID NCT03759288</b>	Crohn's disease	928	<ul style="list-style-type: none"> <li>A 52-Week, Multicenter, Randomized, Double-blind, Placebo and Active-Controlled, Operationally Seamless Phase 2b/3, Parallel group Study to Assess the Efficacy and Safety of Brazikumab in Participants With Moderately to Severely Active Crohn's Disease</li> <li>Stage 1</li> <li>Arm 1: brazikumab high i.v. dose on day 1, 29 and 57 + s.c. brazikumab on day 85 and every 4 weeks through week 48</li> <li>Arm 2: brazikumab low i.v. dose on day 1, 29 and 57 s.c. brazikumab on day 85 and every 4 weeks through week 48</li> <li>Arm 3: placebo</li> <li>Stage 2</li> <li>Arm 1: brazikumab high i.v. dose on day 1, 29 and 57 + s.c. brazikumab on day 85 and every 4 weeks through week 48</li> <li>Arm 2: brazikumab low i.v. dose on day 1, 29 and 57 s.c. brazikumab on day 85 and every 4 weeks through week 48</li> <li>Arm 3: adalimumab s.c. on day 1, 15, 29 and every 2 weeks through week 50</li> </ul>	<ul style="list-style-type: none"> <li>Stage 1 primary endpoint: CDAI remission at week 12</li> <li>Stage 1 secondary endpoints: Endoscopic response at week 12, Clinical remission at week 12, CDAI response at week 12, Response and remission at week 52</li> <li>Stage 2 primary endpoints: Endoscopic response at week 52, Clinical remission at week 52</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2018</li> <li>Data anticipated: &gt;2023</li> </ul>
<b>Phase II EXPEDITION NCT03616821</b>	Ulcerative colitis	256	<ul style="list-style-type: none"> <li>A 54-Week, Multicenter, Randomized, Double-blind, Placebo controlled, Parallel-group Phase 2 Study to Assess the Efficacy and Safety of Brazikumab in Participants with Moderately to Severely Active Ulcerative Colitis</li> <li>Arm 1: brazikumab dose 1 i.v. on day 1, 15 and 43 + s.c. brazikumab from day 71 and every 4 weeks</li> <li>Arm 2: brazikumab dose 2 i.v. on day 1, 15 and 43 + s.c. brazikumab from day 71 and every 4 weeks</li> <li>Arm 3: placebo</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: clinical remission at week 10</li> <li>Secondary endpoint: sustained clinical remission at week 10 and 54</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2018</li> <li>Data anticipated: &gt;2023</li> </ul>



# Brazikumab (IL-23 inhibitor)

## Inflammatory bowel disease (Crohn's disease, ulcerative colitis)

Trial	Population	Patients	Design	Endpoints	Status
<b>Phase II</b> <b>NCT04277546</b>	Ulcerative colitis	165	<ul style="list-style-type: none"> <li>A Phase 2 Open-label, Long-term Extension Safety Study of Brazikumab in Participants with Moderately to Severely Active Ulcerative Colitis</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: safety of long-term treatment with Brazikumab (AEs, clinical laboratory values, vital signs, and ECGs)</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2020</li> <li>Data anticipated: &gt;2023</li> </ul>
<b>Phase I</b> <b>NCT05033431</b>	Healthy volunteers	48	<ul style="list-style-type: none"> <li>Open-label Study to Evaluate the Pharmacokinetics, Safety and Tolerability of a Single Dose of Brazikumab Administered by IV Infusion and SC Injection in Healthy Chinese and White Participants</li> </ul>	<ul style="list-style-type: none"> <li>Primary pharmacokinetic endpoints:               <ul style="list-style-type: none"> <li>C<sub>max</sub></li> <li>AUC<sub>inf</sub></li> <li>AUC<sub>last</sub></li> <li>AUC<sub>0-28d</sub></li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2021</li> <li>Data anticipated: H2 2022</li> </ul>



# PT027 (SABA/ICS, pMDI)

## Asthma

Trial	Population	Patients	Design	Endpoints	Status
<b>Phase III</b> <b>MANDALA</b> <b>NCT03769090</b> <b>Managed by Avillion</b>	Moderate to severe asthma	3132	<ul style="list-style-type: none"> <li>Randomised, double-blind, multicentre, parallel group</li> <li>Treatments (minimum 24-week treatment period)</li> <li>BDA (budesonide albuterol) MDI 80/180 µg prn</li> <li>BDA MDI 160/180 µg prn</li> <li>AS (albuterol sulphate) MDI 180 µg prn</li> <li>Multi-country</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: Time to first severe asthma exacerbation</li> <li>Secondary endpoints: Severe exacerbation rate (annualised); total corticosteroid exposure over the treatment period; asthma Control Questionnaire -5 change from baseline and responder analysis at Week 24; asthma quality of life questionnaire for 12 years and older/paediatric asthma quality of life questionnaire change from baseline and responder analysis at week 24</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2018</li> <li>LPCD: Q1 2021</li> <li>Data readout: Q3 2021</li> <li>Primary endpoint met</li> </ul>
<b>Phase III</b> <b>DENALI</b> <b>NCT03847896</b> <b>Managed by Avillion</b>	Mild to moderate asthma	1001	<ul style="list-style-type: none"> <li>Randomised, double-blind, multicentre and parallel-group</li> <li>Treatments (12 week treatment period)</li> <li>BDA MDI 80/180 µg QID</li> <li>BDA MDI 160/180 µg QID</li> <li>BD MDI 160 µg QID</li> <li>AS MDI 180 µg QID</li> <li>placebo MDI QID</li> <li>Multi-country</li> </ul>	<ul style="list-style-type: none"> <li>Dual primary endpoints: Change from baseline in FEV1 AUC0-6 hours over 12 weeks; change from baseline in trough FEV1 at week 12</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2019</li> <li>LPCD: Q2 2021</li> <li>Data readout: Q3 2021</li> <li>Dual primary endpoints met</li> </ul>
<b>Phase III</b> <b>TYREE</b> <b>NCT04234464</b> <b>Managed by Avillion</b>	Asthma with exercise induced bronchoconstriction	60	<ul style="list-style-type: none"> <li>Randomised, double-blind, multicentre crossover</li> <li>Treatments (single dose)</li> <li>BDA MDI 160/180 µg</li> <li>placebo MDI QID</li> <li>US</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: The maximum percentage fall from post-dose, pre-exercise baseline in forced expiratory volume in 1 second (FEV1) observed up to 60 minutes post-exercise challenge</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2020</li> <li>LPCD: Q3 2020</li> <li>Data readout: Q4 2020</li> <li>Primary endpoint met</li> </ul>



# Tozorakimab (IL-33 ligand mAb)

## Atopic dermatitis, asthma

Trial	Population	Patients	Design	Endpoints	Status
<b>Phase II</b> <b>NCT04212169</b>	Adult subjects with atopic dermatitis	152	<ul style="list-style-type: none"> <li>Randomised, double-blind, placebo-controlled trial</li> <li>Arm 1: tozorakimab s.c.</li> <li>Arm 2: tozorakimab s.c.</li> <li>Arm 3: tozorakimab s.c.</li> <li>Arm 4: placebo s.c.</li> <li>US, Australia, Germany, Poland, UK &amp; Spain</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: change from baseline at week 16 in Eczema Area and Severity Index (EASI) score</li> <li>Secondary endpoints: safety and other efficacy measures</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2019</li> <li>Data anticipated: H2 2022</li> </ul>
<b>Phase II</b> <b>NCT04570657</b>	Adult participants with uncontrolled moderate to severe asthma	228	<ul style="list-style-type: none"> <li>Randomised, double-blind, placebo-controlled trial</li> <li>Arm 1: tozorakimab Dose 1 s.c.</li> <li>Arm 2: tozorakimab Dose 2 s.c.</li> <li>Arm 3: placebo s.c.</li> <li>US, Argentina, Germany, Hungary, Poland, South Africa and UK</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: change from baseline at week 16 in FEV1</li> <li>Secondary endpoints: safety and other efficacy measures</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2020</li> <li>Data anticipated: H2 2022</li> </ul>
<b>Phase I</b> <b>NCT05070312</b>	Healthy subjects	36	<ul style="list-style-type: none"> <li>Randomized, double-blind, placebo-controlled, dose ascending trial</li> <li>Arm 1: tozorakimab Dose 1 s.c.</li> <li>Arm 2: placebo s.c.</li> <li>Arm 3: tozorakimab Dose 2 s.c.</li> <li>Arm 4: placebo s.c.</li> <li>China</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: to characterize the pharmacokinetics of tozorakimab</li> <li>Secondary endpoints: to evaluate the immunogenicity of tozorakimab</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2021</li> <li>LPCD: Q1 2022</li> <li>Data anticipated: H2 2022</li> </ul>



# Tozorakimab (IL-33 ligand mAb)

## COPD

Trial	Population	Patients	Design	Endpoints	Status
<b>Phase III OBERON NCT05166889</b>	Adults with symptomatic COPD with a history of exacerbations	1272	<ul style="list-style-type: none"> <li>Randomised, double-blind, placebo-controlled, parallel group</li> <li>Treatment: 52 week</li> <li>Arm 1: Tozorakimab dose 1 s.c. + standard of care</li> <li>Arm 2: Tozorakimab dose 2 s.c. + standard of care</li> <li>Arm 3: Placebo s.c. + standard of care</li> <li>Global trial: 20 countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: annualized rate of moderate to severe COPD exacerbations (former smokers)</li> <li>Secondary endpoints: annualized rate of moderate to severe COPD exacerbations (former or current smokers), time to moderate to severe COPD exacerbation, change in pre-BD FEV1, E-RS:COPD and SGRQ</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2022</li> <li>Data anticipated: &gt;2023</li> </ul>
<b>Phase III TITANIA NCT05158387</b>	Adults with symptomatic COPD with a history of exacerbations	1272	<ul style="list-style-type: none"> <li>Randomised, double-blind, placebo-controlled, parallel group</li> <li>Treatment: 52 week</li> <li>Arm 1: Tozorakimab dose 1 s.c. + standard of care</li> <li>Arm 2: Tozorakimab dose 2 s.c. + standard of care</li> <li>Arm 3: Placebo s.c. + standard of care</li> <li>Global trial: 19 countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: annualized rate of moderate to severe COPD exacerbations (former smokers)</li> <li>Secondary endpoints: annualized rate of moderate to severe COPD exacerbations (former or current smokers), time to moderate to severe COPD exacerbation, change in pre-BD FEV1, E-RS:COPD and SGRQ</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2022</li> <li>Data anticipated: &gt;2023</li> </ul>
<b>Phase II NCT04631016</b>	Adult subjects with COPD and chronic bronchitis	144	<ul style="list-style-type: none"> <li>Randomised, double-blind, placebo-controlled, parallel group, proof of concept trial</li> <li>Arm 1: Tozorakimab s.c.</li> <li>Arm 2: placebo s.c.</li> <li>US, Australia, Canada, Czech Republic, Denmark, Germany, Hungary, Israel, Netherlands, New Zealand, Poland, South Africa, Spain, Taiwan and UK</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: change from baseline at week 12 in FEV1</li> <li>Secondary endpoints: safety and other efficacy measures</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2021</li> <li>Data anticipated: 2023</li> </ul>



# Evusheld (AZD7442, tixagevimab + cilgavimab)

## Prevention and treatment of COVID-19

Trial	Population	Patients	Design	Endpoints	Status
Phase III PROVENT NCT04625725	Adults having increased risk for inadequate response to active immunisation or having increased risk for SARS-CoV-2 infection	5197	<ul style="list-style-type: none"> <li>Double-blinded, randomised, placebo controlled, multi centre study to determine safety and efficacy in pre-exposure prophylaxis</li> <li>Arm 1: <i>Evusheld</i></li> <li>Arm 2: placebo</li> <li><i>Evusheld</i>/placebo (2:1)</li> <li>USA, UK, Belgium, France, Spain</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: positive symptomatic illness post –dose</li> <li>Secondary endpoints: Incidence of: nucleocapsid antibodies, emergency visits, PCR positive, ADA to <i>Evusheld</i> in serum and <i>Evusheld</i> serum conc.</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2020</li> <li>LPCD: Q1 2021</li> <li>Data readout: Q3 2021</li> <li>Primary endpoint met</li> </ul>
Phase III STORM CHASER NCT04625972	Adults with potential exposure to an identified individual with confirmed SARS-COV2 infection and at risk of developing COVID-19	1121	<ul style="list-style-type: none"> <li>Double-blinded, randomised, placebo controlled, multi centre study to determine safety and efficacy in post-exposure prophylaxis</li> <li>Arm 1: <i>Evusheld</i></li> <li>Arm 2: placebo</li> <li><i>Evusheld</i>/placebo (2:1)</li> <li>USA and UK</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: positive symptomatic illness post –dose</li> <li>Secondary endpoints: Incidence of: nucleocapsid antibodies, COVID-19 related death, all cause mortality, ADA to <i>Evusheld</i> in serum and AZD7442 serum conc.</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2020</li> <li>LPCD: Q1 2021</li> <li>Data readout: Q2 2021</li> <li>Primary endpoint not met</li> </ul>
Phase III TACKLE NCT04723394	Adults with confirmed mild to moderate SARS-COV2 infection. Symptomatic patients with documented positive SARS-Cov-2 molecular test	910	<ul style="list-style-type: none"> <li>Double-blinded, randomised, placebo controlled, multi centre study to determine safety and efficacy for treatment of Covid-19 in non-hospitalised patients</li> <li>Arm 1: <i>Evusheld</i></li> <li>Arm 2: placebo</li> <li><i>Evusheld</i>/placebo (1:1)</li> <li>UK, Germany, Spain, Italy, Hungary, Russia, US, Mexico, Japan, Poland, Czech Republic, Argentina, Brazil, and Ukraine</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: efficacy in the prevention of the composite endpoint of either severe COVID-19 or death from any cause through study day 29</li> <li>Secondary endpoints: A composite of either death from any cause or hospitalisation for COVID-19 complications or sequelae (Day 1 to Day 169). Determine symptom severity and prevention of respiratory failure</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2021</li> <li>LPCD: Q3 2021</li> <li>Data readout: Q4 2021</li> <li>Primary endpoint met</li> </ul>
Phase I NCT04507256	Healthy adults Aged 18-55 years	60	<ul style="list-style-type: none"> <li>Double-blinded, randomised, placebo controlled, single ascending dose trial</li> <li>Arm 1: <i>Evusheld</i></li> <li>Arm 2: placebo</li> <li><i>Evusheld</i>/placebo (10:2)</li> <li>UK</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: safety, tolerability and PK</li> <li>Secondary endpoint: immunogenicity</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2020</li> <li>LPCD: Q3 2020</li> <li>Data readout: Q1 2022</li> </ul>



# Evusheld (AZD7442, tixagevimab + cilgavimab)

## Prevention and treatment of COVID-19

Trial	Population	Patients	Design	Endpoints	Status
Phase I PK Co-formulation NCT05166421	Healthy adults Aged ≥ 18 years	207	<ul style="list-style-type: none"> <li>Open-label, randomised three-arm, single dose trial</li> <li>Arm 1: <i>Evusheld</i> administered as a single co-formulated dose (clonal cell line material)</li> <li>Arm 2: <i>Evusheld</i> administered as two separate doses (clonal cell line material)</li> <li>Arm 3: <i>Evusheld</i> administered as two separate doses (cell pool material)</li> <li><i>Evusheld</i> (1:1:1)</li> <li>USA</li> </ul>	<ul style="list-style-type: none"> <li>Key Endpoints: safety and PK</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2022</li> <li>Data anticipated: 2023</li> </ul>
Phase I TRUST NCT05281601 Partnered	Pediatric participants aged ≥ 29 weeks gestational age to < 18 years at increased risk of developing severe SARS-CoV-2 infection.	100	<ul style="list-style-type: none"> <li>Open-label, single dose, three-cohort trial</li> <li>Cohort 1: Pre-exposure prophylaxis</li> <li>Cohort 2: Mild-to-moderate COVID-19 (cohort yet to open)</li> <li>Cohort 3: Severe COVID-19 (cohort yet to open)</li> <li><i>Evusheld</i></li> <li>USA</li> </ul>	<ul style="list-style-type: none"> <li>Key endpoints: Safety, tolerability and PK</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2022</li> <li>Data anticipated: H2 2022</li> </ul>



# Vaxzevria (SARS-CoV-2)

## Prevention of COVID-19

Approved medicines  
Late-stage development  
Early development

Trial	Population	Patients	Design	Endpoints	Status
Phase IV VICTORIA NCT05057897	Seronegative immunocompromised and immunocompetent individuals who are unvaccinated	360	<ul style="list-style-type: none"> <li>Open-label, non-randomized, multi-cohort, multicenter trial.</li> <li>Vaxzevria i.m.</li> <li>Arm 1: Solid organ transplant</li> <li>Arm 2: Haematopoietic stem cell transplant</li> <li>Arm 3: Solid organ cancer patients receiving cytotoxic therapy</li> <li>Arm 4: Chronic inflammatory disorders</li> <li>Arm 5: Primary immunodeficiency</li> <li>Arm 6: Immunocompetent</li> </ul>	<ul style="list-style-type: none"> <li>Immunogenicity and safety</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2022</li> <li>LPCD: Q3 2022</li> <li>Data anticipated: 2023</li> </ul>
Phase III D8110C00001 (US, global) NCT04516746	Healthy adults Aged 18-65 years	32429	<ul style="list-style-type: none"> <li>Adaptive, double-blinded, randomised placebo-controlled trial</li> <li>Vaxzevria</li> <li>placebo</li> <li>US, Peru, Chile</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: efficacy, safety, tolerability, and reactogenicity</li> <li>Secondary endpoints: immunogenicity</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2020</li> <li>LPCD: Q1 2021</li> <li>Primary data readout Q1 2021</li> </ul>
Phase III COV003 (Brazil) NCT04536051 Partnered	Health professionals and adults with high potential for exposure to SARS-CoV-2 Age 18-55 years	10416	<ul style="list-style-type: none"> <li>Single-blinded, randomised, controlled multicentre trial</li> <li>Vaxzevria</li> <li>Control vaccine: MenACWY</li> <li>Brazil</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: efficacy</li> <li>Secondary endpoints: safety, tolerability, reactogenicity, and immunogenicity</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2020</li> <li>LPCD: Q4 2020</li> </ul>
Phase II/III COV002 (UK) NCT04400838 Partnered	Main efficacy trial: healthy adults aged ≥18 years Healthy adults 56 - <70 years Healthy adults ≥70 years Healthy children 5 – 12 years	10812	<ul style="list-style-type: none"> <li>Single-blinded, randomised, controlled, multicentre trial with sequential age escalation/de-escalation immunogenicity sub-studies that include prime boost</li> <li>Vaxzevria</li> <li>Control vaccine: MenACWY</li> <li>UK</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: efficacy and safety</li> <li>Secondary endpoints: safety, tolerability, reactogenicity, and immunogenicity</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2020</li> <li>LPCD: Q4 2020</li> </ul>
Phase II/III (UK, Brazil, South Africa, Poland) NCT04973449	Healthy adults Aged > 18 years	2590	<ul style="list-style-type: none"> <li>Partially double-blinded, randomized, active-controlled trial in unvaccinated or previously vaccinated participants</li> <li>AZD2816</li> <li>Vaxzevria</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: safety, tolerability</li> <li>Secondary endpoint: immunogenicity</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2021</li> <li>Data readout: Q1 2022</li> <li>Primary endpoint met</li> </ul>

Oncology

CVRM

R&I

Other

V&I

Rare Disease



# Vaxzevria (SARS-CoV-2)

## Prevention of COVID-19

Approved medicines  
Late-stage development  
Early development

Trial	Population	Patients	Design	Endpoints	Status
Phase II COV006 (UK) NCT_JSRCTN15638344 Partnered	Healthy children and adolescents age 6-17 years	261	<ul style="list-style-type: none"> <li>Single-blinded, randomized multicentre trial</li> <li>Vaxzevria</li> <li>Meningococcal Group B vaccine</li> <li>UK</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: safety, tolerability, reactogenicity</li> <li>Secondary endpoints: immunogenicity</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2021</li> <li>LPCD: Q2 2021</li> </ul>
Phase I/II COV001 (UK) NCT04324606 Partnered	Healthy adults Age 18-55 years	1077	<ul style="list-style-type: none"> <li>Single-blinded, randomised, controlled, multicentre trial</li> <li>Vaxzevria</li> <li>Control vaccine: MenACWY</li> <li>UK</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: efficacy and safety</li> <li>Secondary endpoints: safety, tolerability, reactogenicity, and immunogenicity</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2020</li> <li>LPCD: Q2 2020</li> </ul>
Phase I/II COV005 (SA) NCT04444674 Partnered	Healthy adults Age 18-65 years HIV+ subgroup	2130	<ul style="list-style-type: none"> <li>Adaptive, double-blinded, randomised placebo-controlled trial</li> <li>Vaxzevria</li> <li>placebo</li> <li>South Africa</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: efficacy, safety, and immunogenicity</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2020</li> <li>LPCD: Q4 2020</li> </ul>
Phase I/II D8111C00002 (Japan) NCT04568031	Healthy adults Age ≥18 years	256	<ul style="list-style-type: none"> <li>Double-blinded, randomised, placebo-controlled multicentre trial</li> <li>Vaxzevria</li> <li>placebo</li> <li>Japan</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: safety, tolerability, reactogenicity, immunogenicity</li> <li>Secondary endpoints: immunogenicity</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2020</li> <li>LPCD: Q4 2020</li> <li>Data readout: Q4 2020</li> </ul>
Phase I/II COV004 (Kenya) NCT_tbc Partnered	Healthy adults	400	<ul style="list-style-type: none"> <li>Double-blinded, randomised, placebo-controlled multicentre trial</li> <li>Vaxzevria</li> <li>Control vaccine: rabies</li> <li>Kenya</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: safety, tolerability, reactogenicity, immunogenicity</li> <li>Secondary endpoints: immunogenicity</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2020</li> <li>LPCD: Q3 2021</li> </ul>
Phase I COV008 (UK) NCT04816019 Partnered	Healthy adults Age 18-55	54	<ul style="list-style-type: none"> <li>Open label, randomized, dose-escalation trial of Vaxzevria administered intranasally</li> <li>Vaxzevria</li> <li>UK</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: safety, tolerability</li> <li>Secondary endpoint: immunogenicity</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2021</li> <li>LPCD: Q4 2020</li> </ul>



# Nirsevimab (Respiratory syncytial virus mAb-YTE)

## Infection

Approved medicines

Late-stage development

Early development

Trial	Population	Patients	Design	Endpoints	Status
<b>Phase III MELODY NCT03979313</b>	Healthy infants (born 35 weeks 0 days or greater GA)	3000	<ul style="list-style-type: none"> <li>Randomised, Double-blind, placebo-controlled</li> <li>Arm 1: nirsevimab i.m.</li> <li>Arm 2: placebo i.m.</li> <li>Global trial – 31 countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: efficacy</li> <li>Secondary endpoints: safety, PK, ADA</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2021 (safety cohort)</li> <li>LPCD: Q4 2021 (safety cohort)</li> <li>Data Anticipated: H2 2022 (safety cohort)</li> <li>FPCD: Q3 2019 (efficacy cohort)</li> <li>LPCD: Q1 2020 (efficacy cohort)</li> <li>Data readout: Q2 2021 (efficacy cohort)</li> <li>Primary endpoint met</li> </ul>
<b>Phase III CHIMES NCT05110261</b>	Healthy infants (born 29 weeks 0 days or greater GA)	800	<ul style="list-style-type: none"> <li>Randomised, Double-blind, placebo-controlled</li> <li>Arm 1: nirsevimab i.m.</li> <li>Arm 2: placebo i.m.</li> <li>China Only</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: efficacy</li> <li>Secondary endpoints: safety, PK, ADA</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2021</li> <li>Data anticipated: &gt;2023</li> </ul>
<b>Phase IIb NCT02878330</b>	29-35 WK GA (gestational age) infants	1453	<ul style="list-style-type: none"> <li>Randomised, double-blind, placebo-controlled trial</li> <li>Arm 1: nirsevimab i.m.</li> <li>Arm 2: placebo i.m.</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: safety and efficacy</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2016</li> <li>LPCD: Q4 2017</li> <li>Data readout: Q4 2018</li> <li>Primary endpoint met</li> </ul>
<b>Phase II/III MEDLEY NCT03959488</b>	High risk preterm (born 35 weeks 0 day or less GA), CHD and CLD infants eligible to receive Synagis	925	<ul style="list-style-type: none"> <li>Randomised, Double-blind, palivizumab-controlled</li> <li>Arm 1: nirsevimab i.m.</li> <li>Arm 2: Synagis i.m.</li> <li>Global trial – 32 countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: safety and tolerability</li> <li>Secondary endpoints: PK, ADA and descriptive efficacy</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2019</li> <li>LPCD: Q4 2020</li> <li>Data readout: Q2 2021</li> <li>Safety objective met</li> </ul>
<b>Phase II MUSIC NCT04484935</b>	Immunocompromised children who are ≤ 24 months of age at the time of dose administration	100	<ul style="list-style-type: none"> <li>Open-label, Uncontrolled, single-dose trial</li> <li>nirsevimab i.m.</li> <li>Route of administration: i.m.</li> <li>Global trial – 8 countries</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: safety and tolerability</li> <li>Secondary endpoints: PK, ADA, efficacy</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2020</li> <li>Data anticipated: 2023</li> </ul>

Oncology

CVRM

R&I

Other

V&I

Rare Disease



# Nirsevimab (Respiratory syncytial virus mAb-YTE)

## Infection

Approved medicines  
**Late-stage development**  
 Early development

Trial	Population	Patients	Design	Endpoints	Status
Phase I China NCT04840849	Healthy Chinese adults, 18-45 years of age	24	<ul style="list-style-type: none"> <li>Randomised, Double-blind, placebo-controlled</li> <li>Arm 1: nirsevimab i.m.</li> <li>Arm 2: placebo i.m.</li> <li>Route of administration: i.m.</li> <li>China only</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: PK</li> <li>Secondary endpoints: ADA, safety</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2021</li> <li>LPCD: Q2 2021</li> <li>Data anticipated: H1 2022</li> </ul>

Oncology  
 CVRM  
 R&I  
 Other  
**V&I**  
 Rare Disease



# BioPharmaceuticals – early-stage development



# AZD2373

## Chronic kidney disease

Approved medicines

Late-stage development

Early development

Oncology

CVRM

R&I

Other

V&I

Rare Disease

Trial	Population	Patients	Design	Endpoints	Status
Phase I NCT04269031	Healthy volunteers	30	<ul style="list-style-type: none"><li>SAD. Dose escalation in 6 cohorts with 6 volunteers receiving AZD2373 and 2 volunteers receiving placebo in each cohort</li><li>Arm 1: AZD2373 s.c.</li><li>Arm 2: placebo s.c.</li><li>US</li></ul>	<ul style="list-style-type: none"><li>Primary endpoints: safety and tolerability</li><li>Secondary endpoint: PK parameters</li></ul>	<ul style="list-style-type: none"><li>FPCD: Q1 2020</li><li>LPCD: Q3 2021</li><li>Data anticipated: H1 2022</li></ul>



# AZD2693 (antisense oligonucleotide)

## NASH

Trial	Population	Patients	Design	Endpoints	Status
Phase I NCT04142424	Healthy volunteers	62	<ul style="list-style-type: none"><li>SAD. 6 cohorts with 6 volunteers receiving AZD2693 and 2 volunteers receiving placebo in each cohort</li><li>Arm 1: AZD2693 s.c.</li><li>Arm 2: placebo s.c.</li><li>US</li></ul>	<ul style="list-style-type: none"><li>Primary endpoints: safety and tolerability</li><li>Secondary endpoint: PK</li></ul>	<ul style="list-style-type: none"><li>FPCD: Q4 2019</li><li>LPCD: Q3 2021</li><li>Data readout: Q4 2021</li></ul>
Phase I NCT04142424	NASH/NAFLD F0-F3	60	<ul style="list-style-type: none"><li>MAD. 3 cohorts receiving AZD2693 and placebo in each cohort</li><li>Arm 1: AZD2693 s.c.</li><li>Arm 2: placebo s.c.</li><li>US</li></ul>	<ul style="list-style-type: none"><li>Primary endpoints: safety and tolerability</li><li>Secondary endpoint: PK</li></ul>	<ul style="list-style-type: none"><li>FPCD: Q2 2021</li><li>Data anticipated: H1 2022</li></ul>



# AZD3366

## Cardiovascular disease

Approved medicines

Late-stage development

Early development

Oncology

CVRM

R&I

Other

V&I

Rare Disease

Trial	Population	Patients	Design	Endpoints	Status
Phase I NCT04588727	Healthy volunteers	103	<ul style="list-style-type: none"><li>• SAD trial</li><li>• Part A</li><li>• 7 cohorts with 6 volunteers receiving AZD3366 and 2 volunteers receiving placebo in each cohort; three cohorts with Japanese subjects 5 receiving AZD3366</li><li>• 1 receiving placebo; 1 Chinese cohort of 6 receiving AZD3366 2 receiving placebo</li><li>• Arm 1: AZD3366</li><li>• Arm 2: placebo</li><li>• Part B</li><li>• 12 subjects</li><li>• Arm 1: AZD3366 + <i>Brilinta</i> + ASA</li><li>• Arm 2: placebo + <i>Brilinta</i> + ASA</li></ul>	<ul style="list-style-type: none"><li>• Primary endpoints: safety and tolerability</li><li>• Secondary endpoint: PK parameters</li></ul>	<ul style="list-style-type: none"><li>• FPCD: Q4 2020</li><li>• Data anticipated: H1 2022</li></ul>



# AZD3427 (relaxin)

## Heart failure

Trial	Population	Patients	Design	Endpoints	Status
Phase I NCT04630067	SAD – Healthy volunteers MAD – Heart failure	104	<ul style="list-style-type: none"><li>• Multicentre single and multiple ascending dose study</li><li>• Part A SAD 6 cohorts</li><li>• Arm 1: AZD3427</li><li>• Arm 2: placebo</li><li>• Part B MAD</li><li>• Arm 1: AZD3427</li><li>• Arm 2: placebo</li><li>• US</li></ul>	<ul style="list-style-type: none"><li>• Primary endpoints: safety and tolerability</li></ul>	<ul style="list-style-type: none"><li>• FPCD: Q4 2020</li><li>• LPCD: Q3 2022</li><li>• Data anticipated: H2 2022</li></ul>



# AZD4831 (MPO inhibitor)

## Cardiovascular disease

Approved medicines

Late-stage development

Early development

Oncology

CVRM

R&I

Other

V&I

Rare Disease

Trial	Population	Patients	Design	Endpoints	Status
Phase I NCT05236543	Healthy Patients	14	<ul style="list-style-type: none"><li>• Open label</li><li>• AZD4831</li><li>• AZD4831 and Itraconazole</li><li>• UK</li></ul>	<ul style="list-style-type: none"><li>• Primary endpoints: PK parameters</li><li>• Secondary endpoints: safety and tolerability</li></ul>	<ul style="list-style-type: none"><li>• FPCD: Q1 2022</li><li>• LPCD: Q2 2022</li><li>• Data anticipated: 2023</li></ul>



# AZD5462 (relaxin)

## Heart failure

Approved medicines

Late-stage development

Early development

Oncology

CVRM

R&I

Other

V&I

Rare Disease

Trial	Population	Patients	Design	Endpoints	Status
Phase I NCT04994106	SAD – Healthy volunteers MAD – Healthy volunteers	0	<ul style="list-style-type: none"><li>• Single centre single and multiple ascending dose study</li><li>• Part A SAD 8 cohorts</li><li>• Arm 1: AZD5462</li><li>• Arm 2: placebo</li><li>• Part B MAD 5 cohorts</li><li>• Arm 1: AZD5462</li><li>• Arm 2: placebo</li><li>• US</li></ul>	<ul style="list-style-type: none"><li>• Primary endpoints: safety and tolerability</li></ul>	<ul style="list-style-type: none"><li>• FPCD: Q4 2021</li><li>• Data anticipated: H2 2022</li></ul>



# AZD5718 (FLAP inhibitor)

## Cardiovascular disease & Chronic Kidney Disease

Trial	Population	Patients	Design	Endpoints	Status
<b>Phase IIb FLAIR NCT04492722</b>	Proteinuric CKD	632	<ul style="list-style-type: none"> <li>Randomised, double-blind, placebo-controlled, multicentre, dose-ranging trial</li> <li>AZD5718</li> <li>placebo</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: dose-response efficacy, safety, PK</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2020</li> <li>Data anticipated: 2023</li> </ul>
<b>Phase IIa NCT03317002</b>	CAD	129	<ul style="list-style-type: none"> <li>Arm 1: AZD5718 Dose A</li> <li>Arm 2: AZD5718 Dose B</li> <li>Arm 3: placebo</li> <li>Global trial – three countries in Europe</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: PD effect of AZD5718 by assessment of u-LTE4</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2017</li> <li>LPCD: Q4 2019</li> <li>Data readout: Q1 2021</li> </ul>
<b>Phase I NCT03948451</b>	Healthy patients	6	<ul style="list-style-type: none"> <li>Open label hADME trial</li> <li>a single oral dose of 14C-AZD5718</li> <li>UK</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: Mass balance, with routes and rates of elimination of 14C-AZD5718; Metabolite profiling and structural identification; PK and total radioactivity</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2019</li> <li>LPCD: Q2 2019</li> <li>Data readout: Q4 2019</li> </ul>
<b>Phase I NCT04087187</b>	Healthy patients	14	<ul style="list-style-type: none"> <li>BA trial. Open-label, randomised, 3-period, 3-treatment, crossover design</li> <li>AZD5718</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: safety and tolerability, PK</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2019</li> <li>LPCD: Q4 2019</li> <li>Data readout: Q1 2020</li> </ul>
<b>Phase I NCT04210388</b>	Healthy patients	12	<ul style="list-style-type: none"> <li>BA trial. Open-label, randomised, single-dose, combined 2x2 dose and 3x3 dose crossover design in fixed sequence.</li> <li>AZD5718</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: bioavailability, safety and tolerability</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2020</li> <li>LPCD: Q1 2020</li> <li>Data readout: Q2 2020</li> </ul>
<b>Phase I NCT04734275</b>	Healthy patients	16	<ul style="list-style-type: none"> <li>BA trial. Open-label, randomised, single-dose, 3-period, single dose, crossover design</li> <li>AZD5718</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: bioavailability, safety and tolerability</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2021</li> <li>LPCD: Q1 2021</li> <li>Data readout: Q4 2021</li> </ul>



# AZD7503 (antisense oligonucleotide)

## NASH

Approved medicines  
Late-stage development  
Early development

Trial	Population	Patients	Design	Endpoints	Status
Phase I NCT05143905	Healthy volunteers	64	<ul style="list-style-type: none"><li>SAD. 6 cohorts with 6 volunteers receiving AZD7503 and 2 volunteers receiving placebo in each cohort</li><li>Arm 1: AZD7503 s.c.</li><li>Arm 2: placebo s.c.</li><li>US</li></ul>	<ul style="list-style-type: none"><li>Primary endpoints: safety and tolerability</li><li>Secondary endpoint: PK</li></ul>	<ul style="list-style-type: none"><li>FPCD: Q4 2021</li><li>Data anticipated: 2023</li></ul>

Oncology

CVRM

R&I

Other

V&I

Rare Disease



# AZD8233 (PCSK9 inhibitor, subcutaneous)

## Dyslipidaemia

Trial	Population	Patients	Design	Endpoints	Status
Phase II SOLANO NCT04964557	Dyslipidaemia	376	<ul style="list-style-type: none"> <li>Arm 1: AZD8233 dose</li> <li>Arm 2: placebo</li> <li>28 weeks</li> <li>8 countries (US, Slovakia, Denmark, Poland, Spain, Netherlands, Czech Republic, Hungary)</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: safety</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2021</li> <li>LPCD: Q4 2021</li> <li>Data anticipated: H2 2022</li> </ul>
Phase II ETESIAN NCT04641299	Dyslipidaemia	108	<ul style="list-style-type: none"> <li>Arm 1: High AZD8233 dose</li> <li>Arm 2: Medium AZD8233 dose</li> <li>Arm 3: Low AZD8233 dose</li> <li>Arm 4: placebo</li> <li>12 weeks</li> <li>3 countries (US, Slovakia and Denmark)</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: efficacy</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2020</li> <li>LPCD: Q1 2021</li> <li>Data readout: Q3 2021</li> </ul>
Phase II HAYATE NCT04823611	Dyslipidaemia	82	<ul style="list-style-type: none"> <li>Part A: 11 subjects randomized in an 8:3 ratio</li> <li>Arm 1: AZD8233 dose</li> <li>Arm 2: placebo</li> <li>Part B: 60 subjects randomized across three different treatment arms in a 1:1:1 ratio</li> <li>Arm 1: Medium AZD8233 dose</li> <li>Arm 2: Low AZD8233 dose</li> <li>Arm 3: placebo</li> <li>Part C: 11 subjects randomized in an 8:3 ratio</li> <li>Arm 1: Medium AZD8233 dose</li> <li>Arm 2: placebo</li> <li>12 weeks</li> <li>Japan</li> </ul>	<ul style="list-style-type: none"> <li>Part A primary endpoints: safety and tolerability</li> <li>Part B primary endpoint: efficacy</li> <li>Part C primary endpoints: safety and tolerability</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2021</li> <li>LPCD: Q1 2022</li> <li>Data anticipated: H2 2022</li> </ul>
Phase I NCT03593785	Healthy subjects	72	<ul style="list-style-type: none"> <li>SAD trial</li> <li>7 cohorts</li> <li>Arm1: AZD8233</li> <li>Arm 2: placebo</li> <li>US</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: safety and tolerability</li> <li>Secondary endpoints: PK and PD parameters</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2018</li> <li>LPCD: Q3 2019</li> <li>Data readout: Q2 2021</li> </ul>
Phase I NCT04155645	Dyslipidaemia	33	<ul style="list-style-type: none"> <li>MAD trial</li> <li>3 cohorts</li> <li>Arm1: AZD8233</li> <li>Arm 2: placebo</li> <li>US</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: safety and tolerability</li> <li>Secondary endpoints: PK and PD parameters</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2020</li> <li>LPCD: Q4 2020</li> <li>Data readout: Q3 2021</li> </ul>



# AZD8601 (VEGF-A modified RNA)

## Cardiovascular disease

Approved medicines

Late-stage development

Early development

Oncology

CVRM

R&I

Other

V&I

Rare Disease

Trial	Population	Patients	Design	Endpoints	Status
Phase IIa NCT03370887	HF	33	<ul style="list-style-type: none"><li>Arm 1: AZD8601 Dose A</li><li>Arm 2: AZD8601 Dose B</li><li>Arm 3: placebo</li><li>Finland, Germany</li></ul>	<ul style="list-style-type: none"><li>Primary endpoints: safety and tolerability</li></ul>	<ul style="list-style-type: none"><li>FPCD: Q1 2018</li><li>LPCD: Q1 2021</li><li>Data readout: Q3 2021</li></ul>



# AZD9977 (MR modulator)

## Heart failure

Approved medicines

Late-stage development

Early development

Trial	Population	Patients	Design	Endpoints	Status
<b>Phase IIb MIRACLE NCT04595370</b>	Heart failure with chronic kidney disease	500	<ul style="list-style-type: none"> <li>Randomised, stratified according to T2DM and eGFR (<math>\geq 20</math> to <math>&lt; 30</math> mL/min / <math>\geq 30</math> to <math>&lt; 45</math> mL/min / <math>\geq 45</math> mL/min) for 12 weeks:</li> <li>Arm 1: AZD9977 A + <i>Farxiga</i> 10 mg</li> <li>Arm 2: AZD9977 B + <i>Farxiga</i> 10 mg</li> <li>Arm 3: AZD9977 C + <i>Farxiga</i> 10 mg</li> <li>Arm 4: <i>Farxiga</i> 10 mg</li> <li>12 weeks</li> <li>Trial conducted in 19 countries globally</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: percent change from baseline in UACR at 12 weeks</li> <li>Secondary endpoints: percent change from baseline in UACR at 12 weeks to assess dose-response relationship; dose-response relationship of <i>Farxiga</i> and 3 doses of AZD9977 combined with <i>Farxiga</i> on UACR; safety, tolerability and serum potassium values; eGFR</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2021</li> <li>Data anticipated: 2023</li> </ul>
<b>Phase I NCT04469907</b>	Renal impairment	18	<ul style="list-style-type: none"> <li>AZD9977</li> <li>US</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: PK parameters</li> <li>Secondary endpoints: safety and tolerability</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2020</li> <li>LPCD: Q4 2021</li> <li>Data anticipated: H1 2022</li> </ul>
<b>Phase I NCT04686591</b>	Healthy volunteers	8	<ul style="list-style-type: none"> <li>hADME trial</li> <li>single oral dose 14C-AZD9977</li> <li>UK</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: absolute bioavailability, the mass balance, rates and routes of elimination</li> <li>Secondary endpoints: safety and tolerability</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2021</li> <li>LPCD: Q1 2021</li> <li>Data anticipated: H1 2022</li> </ul>
<b>Phase I NCT04798222</b>	Healthy volunteers	20	<ul style="list-style-type: none"> <li>Rel BA trial</li> <li>Study to Assess the Bioavailability of Different Formulations of AZD9977 and Dapagliflozin and Influence of Food in Selected Formulations</li> <li>Germany</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: relative bioavailability of 9977 and Dapagliflozin in different capsule formulations; PK parameters</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2021</li> <li>LPCD: Q3 2021</li> <li>Data anticipated: H1 2022</li> </ul>

Oncology

CVRM

R&I

Other

V&I

Rare Disease



# Cotadutide (GLP-1-glucagon agonist)

## Diabetes/CKD, NASH

Trial	Population	Patients	Design	Endpoints	Status
<b>Phase II NCT04019561</b>	Obese patients with non-alcoholic fatty liver disease (NAFLD)/non-alcoholic steatohepatitis (NASH)	74	<ul style="list-style-type: none"> <li>• Arm 1: cotadutide high dose s.c.</li> <li>• Arm 2: placebo high dose s.c.</li> <li>• Arm 3: cotadutide low dose s.c.</li> <li>• Arm 4: placebo low dose s.c.</li> <li>• US</li> </ul>	<ul style="list-style-type: none"> <li>• Primary endpoints: safety and tolerability</li> <li>• Secondary endpoints: change in hepatic fat fraction; change in liver fat volume; change in visceral adipose tissue</li> </ul>	<ul style="list-style-type: none"> <li>• FPCD: Q4 2019</li> <li>• LPCD: Q2 2021</li> <li>• Data anticipated: H1 2022</li> </ul>
<b>Phase II NCT04515849</b>	Chronic kidney disease with type 2 diabetes mellitus	225	<ul style="list-style-type: none"> <li>• Arm 1: cotadutide 100 micrograms</li> <li>• Arm 2: cotadutide 300 micrograms</li> <li>• Arm 3: cotadutide 600 micrograms</li> <li>• Arm 4: semaglutide</li> <li>• Arm 5: placebo</li> </ul>	<ul style="list-style-type: none"> <li>• Primary endpoint: efficacy change in UACR</li> <li>• Secondary endpoints: Change in HbA1c; change in glucose measured by CGM; effects on body weight; safety, tolerability, Immunogenicity</li> </ul>	<ul style="list-style-type: none"> <li>• FPCD: Q4 2020</li> <li>• LPCD: Q3 2021</li> <li>• Data anticipated: H1 2022</li> </ul>



# MEDI6570

## Cardiovascular

Approved medicines

Late-stage development

Early development

Oncology

CVRM

R&I

Other

V&I

Rare Disease

Trial	Population	Patients	Design	Endpoints	Status
Phase IIb NCT04610892	Post MI	400	<ul style="list-style-type: none"> <li>Evaluation of anti-inflammatory potential and effect on surrogates for atherosclerotic and heart failure (HF) events</li> <li>Arm 1: High MEDI6570 dose</li> <li>Arm 2: Medium MEDI6570 dose</li> <li>Arm 3: Low MEDI6570 dose</li> <li>Arm 4: placebo</li> <li>US, Canada, Hungary, Japan, Czech Republic, Italy, Spain, Netherlands, Poland, UK, Australia, Russia</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: efficacy and safety</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2020</li> <li>Data anticipated: 2023</li> </ul>



# MEDI8367

## Chronic kidney disease

Approved medicines

Late-stage development

Early development

Oncology

CVRM

R&I

Other

V&I

Rare Disease

Trial	Population	Patients	Design	Endpoints	Status
Phase I NCT04365218	Healthy volunteers CKD	70	<ul style="list-style-type: none"><li>SAD trial</li><li>6 cohorts</li><li>Arm 1: MEDI8367 s.c.</li><li>Arm 2: placebo s.c.</li><li>US</li></ul>	<ul style="list-style-type: none"><li>Primary endpoints: safety and tolerability</li><li>Secondary endpoints: PK parameters, ADA</li></ul>	<ul style="list-style-type: none"><li>FPCD: Q3 2020</li><li>LPCD: Q4 2020</li><li>Data anticipated: H1 2022</li></ul>



# Tozorakimab (IL-33 ligand mAb)

## Diabetic kidney disease

Approved medicines

Late-stage development

Early development

Oncology

CVRM

R&I

Other

V&I

Rare Disease

Trial	Population	Patients	Design	Endpoints	Status
Phase II NCT04170543	Adult patients with diabetic kidney disease	565	<ul style="list-style-type: none"><li>• Arm A: tozorakimab dose 1 + <i>Farxiga</i></li><li>• Arm B: tozorakimab dose 2 + <i>Farxiga</i></li><li>• Arm C: tozorakimab dose 3 + <i>Farxiga</i></li><li>• Arm D: tozorakimab dose 4 + <i>Farxiga</i></li><li>• Arm E: placebo + <i>Farxiga</i></li><li>• USA, Canada, Japan and additional countries.</li></ul>	<ul style="list-style-type: none"><li>• Primary endpoint: Urine albumin creatinine ratio (UACR)</li><li>• Secondary endpoints: Safety and other efficacy measures</li></ul>	<ul style="list-style-type: none"><li>• FPCD: Q4 2019</li><li>• Data anticipated: 2023</li></ul>



# Zibotentan (endothelin receptor antagonist)

## Chronic kidney disease

Trial	Population	Patients	Design	Endpoints	Status
Phase IIb ZENITH-CKD NCT04724837	CKD	660	<ul style="list-style-type: none"> <li>Part A: 132 participants equally randomised across 4 arms:               <ul style="list-style-type: none"> <li>Arm 1: zibotentan dose A + <i>Farxiga</i> 10 mg once daily</li> <li>Arm 2: zibotentan dose A once daily</li> <li>Arm 3: <i>Farxiga</i> 10 mg once daily</li> <li>Arm 4: placebo once daily</li> </ul> </li> <li>Part B: 528 participants equally randomised across 6 arms:               <ul style="list-style-type: none"> <li>Arm 1: zibotentan dose C + <i>Farxiga</i> 10 mg once daily</li> <li>Arm 2: zibotentan dose B + <i>Farxiga</i> 10 mg once daily</li> <li>Arm 3: zibotentan dose A + <i>Farxiga</i> 10 mg once daily</li> <li>Arm 4: zibotentan dose A once daily</li> <li>Arm 5: <i>Farxiga</i> 10 mg once daily</li> <li>Arm 6: placebo once daily</li> </ul> </li> <li>Global</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: change in log-transformed UACR from baseline to week 12.</li> <li>Secondary endpoints: change in log-transformed UACR from baseline to week 12; change in blood pressure from baseline (Visit 2) to week 12; least squares mean change of UACR at week 12 from the 3 Zibo/Dapa dose groups and the <i>Farxiga</i> monotherapy group; change in eGFR from baseline to week 1, week 12 and week 14; change in eGFR from week 1 to week 12.</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2021</li> <li>Data anticipated: H2 2022</li> </ul>
Phase I NCT04991571	Healthy volunteers	28	<ul style="list-style-type: none"> <li>Part 1 of the study is intended to collect samples for Metabolites in Safety Testing (MIST) analysis after administration of multiple doses of zibotentan.</li> <li>Part 2 of the study is designed to evaluate the relative bioavailability of zibotentan and dapagliflozin after dosing with two different fixed-dose combination (FDC) formulations and dosing with separate formulations of zibotentan and dapagliflozin</li> </ul>	<ul style="list-style-type: none"> <li>Part 1: Metabolites in Safety Testing MIST analysis. No PK statistical analysis will be performed.</li> <li>Part 2: PK parameters</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2021</li> <li>LPCD: Q4 2021</li> <li>Data anticipated: H1 2022</li> </ul>



# Zibotentan (endothelin receptor antagonist)

## Liver Cirrhosis with Features of portal hypertension

Trial	Population	Patients	Design	Endpoints	Status
Phase I NCT05112419	Cohort 1: Moderate hepatic impairment/moderate renal impairment patients Cohort 2: Healthy volunteers	24	<ul style="list-style-type: none"> <li>Single dose, single centre, Open label, Parallel group study to investigate the pharmacokinetics, safety and tolerability of 5 mg Zibotentan</li> </ul>	<ul style="list-style-type: none"> <li>Primary Endpoints:</li> <li>Area under plasma concentration-time curve from time zero to infinity (AUC<sub>inf</sub>)</li> <li>Area under the plasma concentration-curve from time zero to time of last quantifiable concentration (AUC<sub>last</sub>)</li> <li>Maximum observed plasma concentration (C<sub>max</sub>)</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2021</li> <li>LPCD: Q4 2021</li> <li>Data anticipated: H1 2022</li> </ul>



# AZD1402 (IL-4 receptor alpha antagonist)

## Asthma

Trial	Population	Patients	Design	Endpoints	Status
<b>Phase IIa</b> <b>Aputura</b> <b>NCT04643158</b>	Patients with asthma on medium dose inhaled corticosteroids	345	<ul style="list-style-type: none"> <li>Randomised, placebo-controlled, double-blinded, multicentre, 2-part trial.</li> <li>Part 1 population with asthma controlled on medium dose ICS-LABA               <ul style="list-style-type: none"> <li>Part 1a                   <ul style="list-style-type: none"> <li>Arm 1: AZD1402 Dose 1 (low) (DPI)</li> <li>Arm 2: AZD1402 Dose 2 (DPI)</li> <li>Arm 3: placebo</li> </ul> </li> <li>Part 1b                   <ul style="list-style-type: none"> <li>Arm 1: AZD1402 Dose 3 (high) (DPI)</li> <li>Arm 2: placebo</li> </ul> </li> </ul> </li> <li>Part 2 population uncontrolled on medium dose ICS-LABA               <ul style="list-style-type: none"> <li>Arm 1: AZD1402 Dose 1 (DPI)</li> <li>Arm 2: AZD1402 Dose 2 (DPI)</li> <li>Arm 3: AZD1402 Dose 3 (DPI)</li> <li>Arm 4: placebo</li> </ul> </li> <li>Ukraine, Australia, Germany, Hungary, Korea, Poland, Spain, UK</li> </ul>	<ul style="list-style-type: none"> <li>Part 1 primary endpoints: safety and tolerability, PK</li> <li>Part 2 primary endpoint: change in FEV1</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2021</li> <li>Data anticipated: H2 2022</li> </ul>



# AZD4604 (inhaled JAK-1 inhibitor)

## Asthma

Approved medicines

Late-stage development

Early development

Trial	Population	Patients	Design	Endpoints	Status
<b>Phase I</b> <b>NCT04769869</b>	Healthy subjects and patients with mild asthma	137	<ul style="list-style-type: none"><li>• SAD/MAD/POM trial</li><li>• Part 1 SAD</li><li>• Arm 1: AZD0449 (DPI)</li><li>• Arm 2: placebo (DPI)</li><li>• Part 2 MAD</li><li>• Arm 1: AZD0449 (DPI)</li><li>• Arm 2: placebo (DPI)</li><li>• Part 3 POM</li><li>• Arm 1: AZD0449 (DPI)</li><li>• Arm 2: placebo (DPI)</li><li>• UK</li></ul>	<ul style="list-style-type: none"><li>• Primary endpoints: safety and tolerability</li><li>• Secondary endpoints: PK parameters, FENO</li></ul>	<ul style="list-style-type: none"><li>• FPCD: Q4 2021</li><li>• Data anticipated: 2023</li></ul>

Oncology

CVRM

R&I

Other

V&I

Rare Disease



# AZD5055 (oral porcupine inhibitor)

## Idiopathic pulmonary fibrosis (IPF) and other ILDs with progressive fibrosis

Trial	Population	Patients	Design	Endpoints	Status
Phase I NCT05134727	Healthy subjects	104	<ul style="list-style-type: none"> <li>SAD/MAD trial</li> <li>Part 1a (fasted state) and 1b (fed state) SAD</li> <li>Arm 1: AZD5055 (oral suspension)</li> <li>Arm 2: Placebo (oral suspension)</li> <li>Part 2 MAD</li> <li>Arm 1: AZD5055 (oral suspension)</li> <li>Arm 2: Placebo (oral suspension)</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: safety and tolerability</li> <li>Secondary endpoints: PK parameters, food effect</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2021</li> <li>Data anticipated: 2023</li> </ul>



# AZD8630 (inhaled TSLP)

## Asthma

Approved medicines

Late-stage development

Early development

Trial	Population	Patients	Design	Endpoints	Status
Phase I NCT05110976 Partnered	Healthy subjects and patients with asthma	232	<ul style="list-style-type: none"><li>SAD &amp; MAD trial</li></ul>	<ul style="list-style-type: none"><li>Primary endpoints: safety and tolerability</li><li>Secondary endpoints: PK parameters, FENO</li></ul>	<ul style="list-style-type: none"><li>FPCD: Q4 2021</li><li>Data anticipated: 2023</li></ul>

Oncology

CVRM

R&I

Other

V&I

Rare Disease



# AZD4041 (orexin 1 receptor antagonist)

## Opioid use disorder

Trial	Population	Patients	Design	Endpoints	Status
<b>Phase I</b> <b>NCT04076540</b> Partnered with Eolas Therapeutics Inc and NIH	Healthy volunteers	48	<ul style="list-style-type: none"> <li>Randomised, double blind, SAD trial</li> <li>Arm 1: AZD4041</li> <li>Arm 2: placebo</li> <li>US only</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: safety and tolerability</li> <li>Secondary endpoints: PK, PD</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2019</li> <li>Data readout: Q4 2021</li> <li>Primary endpoint met</li> </ul>
<b>Phase I</b> <b>NCT05209334</b> Partnered with Eolas Therapeutics Inc and NIH	Healthy volunteers	36	<ul style="list-style-type: none"> <li>Randomised, double blind MAD trial</li> <li>Arm 1: AZD4041</li> <li>Arm 2: placebo</li> <li>Canada only</li> </ul>	<ul style="list-style-type: none"> <li>Primary Endpoints: safety and tolerability</li> <li>Secondary endpoint: PK</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2022</li> <li>LPCD: Q2 2022</li> <li>Data anticipated: H2 2022</li> </ul>



# MEDI0618 (PAR2 antagonist mAb)

## Osteoarthritis pain

Trial	Population	Patients	Design	Endpoints	Status
Phase I NCT04198558	Healthy volunteers	64	<ul style="list-style-type: none"><li>• SAD trial</li><li>• Arm 1: MEDI0618 i.v.</li><li>• Arm 2: placebo i.v.</li><li>• Arm 3: MEDI0618 s.c</li><li>• Arm 4: placebo s.c</li><li>• Europe only</li></ul>	<ul style="list-style-type: none"><li>• Primary endpoints: safety and tolerability</li><li>• Secondary endpoint: PK</li></ul>	<ul style="list-style-type: none"><li>• FPCD: Q4 2019</li><li>• Data anticipated: H1 2022</li></ul>



# MEDI1341 (alpha-synuclein mAb)

## Parkinson's disease

Approved medicines  
Late-stage development  
Early development

Trial	Population	Patients	Design	Endpoints	Status
Phase I NCT03272165 Partnered	Healthy volunteers	48	<ul style="list-style-type: none"><li>SAD trial</li><li>Arm 1: MEDI1341 i.v.</li><li>Arm 2: placebo i.v.</li><li>US only</li></ul>	<ul style="list-style-type: none"><li>Primary endpoints: safety and tolerability</li><li>Secondary endpoints: PK, PD</li></ul>	<ul style="list-style-type: none"><li>FPCD: Q4 2017</li><li>LPCD: Q4 2020</li><li>Data readout: Q4 2021</li></ul>
Phase I NCT04449484 Partnered	Parkinson's Disease	25	<ul style="list-style-type: none"><li>MAD trial</li><li>Arm 1: MEDI1341 i.v.</li><li>Arm 2: placebo i.v.</li><li>US only</li></ul>	<ul style="list-style-type: none"><li>Primary endpoints: safety and tolerability</li><li>Secondary endpoints: PK, PD</li></ul>	<ul style="list-style-type: none"><li>FPCD: Q3 2020</li><li>LPCD: Q3 2021</li><li>Data anticipated: H2 2022</li></ul>

Oncology

CVRM

R&I

Other

V&I

Rare Disease



# MEDI7352 (NGF TNF bispecific mAb)

## Osteoarthritis pain

Trial	Population	Patients	Design	Endpoints	Status
<b>Phase IIb</b> NCT04675034	Painful osteoarthritis of the knee	300	<ul style="list-style-type: none"> <li>MAD trial</li> <li>Arm 1: MEDI7352 s.c</li> <li>Arm 2: placebo s.c</li> <li>Global (7 countries)</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: dose response</li> <li>Secondary endpoints: safety, tolerability, PK, PD, ADA</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2021</li> <li>Data anticipated: 2023</li> </ul>
<b>Phase IIa</b> NCT03755934	Painful diabetic neuropathy	271	<ul style="list-style-type: none"> <li>MAD trial</li> <li>Arm 1: MEDI7352 i.v.</li> <li>Arm 2: placebo i.v.</li> <li>Europe only</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: dose response</li> <li>Secondary endpoints: safety, tolerability, PK, PD</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2018</li> <li>Data anticipated: 2023</li> </ul>
<b>Phase I</b> NCT02508155	Painful osteoarthritis of the knee	160	<ul style="list-style-type: none"> <li>SAD &amp; MAD trial</li> <li>Arm 1: MEDI7352 i.v.</li> <li>Arm 2: placebo i.v.</li> <li>Arm 3: MEDI7352 s.c</li> <li>Arm 4: placebo s.c</li> <li>Europe only</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: safety and tolerability</li> <li>Secondary endpoints: PK, PD</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2016</li> <li>LPCD: Q4 2020</li> <li>Data readout: Q2 2021</li> </ul>
<b>Phase I</b> NCT04770428	Healthy volunteers Japanese and Caucasian	20	<ul style="list-style-type: none"> <li>MAD trial</li> <li>Arm 1: MEDI7352 s.c</li> <li>Arm 2: placebo s.c</li> <li>Europe only</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: safety and tolerability</li> <li>Secondary endpoints: PK, PD, ADA</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q2 2021</li> <li>LPCD: Q3 2021</li> <li>Data readout: Q4 2021</li> </ul>



Rare Disease – approved  
medicines and late-stage  
pipeline



# Soliris (anti-complement C5 mAb)

## Neurology

Approved medicines

Late-stage development

Early development

Trial	Population	Patients	Design	Endpoints	Status
Phase III ECU-GBS-301 NCT04752566	Guillain-Barré syndrome	57	<ul style="list-style-type: none"><li>• Arm 1: <i>Soliris</i> i.v. once weekly for 4 weeks</li><li>• Arm 2: Placebo</li><li>• Japan-only</li></ul>	<ul style="list-style-type: none"><li>• Primary endpoint: Time to first reaching a Hughes functional grading scale score <math>\leq 1</math></li></ul>	<ul style="list-style-type: none"><li>• FPCD: Q1 2021</li><li>• Data anticipated: 2023</li></ul>

Oncology

CVRM

R&I

Other

V&I

Rare Disease



# Ultomiris (anti-complement C5 mAb)

## Haematology & nephrology

Trial	Population	Patients	Design	Endpoints	Status
Phase III ALXN-1210-PNH-303 NCT03748823	PNH and aHUS	136	<ul style="list-style-type: none"> <li>• <i>Ultomiris</i> s.c.</li> </ul>	<ul style="list-style-type: none"> <li>• Primary endpoint: Day 71 serum <i>Ultomiris</i> C trough</li> </ul>	<ul style="list-style-type: none"> <li>• FPCD: Q1 2019</li> <li>• Data readout: Q2 2020</li> <li>• Primary endpoint met</li> </ul>
Phase III ALXN-1210-TMA-315 NCT04743804	Thrombotic microangiopathy associated with a trigger	100	<ul style="list-style-type: none"> <li>• Arm 1: <i>Ultomiris</i> Q8W</li> <li>• Arm 2: Placebo</li> </ul>	<ul style="list-style-type: none"> <li>• Primary endpoint: Complete TMA response</li> <li>• Secondary endpoints: Time to complete TMA response, Hematologic response, Renal response</li> </ul>	<ul style="list-style-type: none"> <li>• FPCD: Q3 2021</li> <li>• Data anticipated: &gt;2023</li> </ul>
Phase III ALXN-1210-TM-313 NCT04543591	Thrombotic microangiopathy associated with haematopoietic stem cell transplant	184	<ul style="list-style-type: none"> <li>• Arm 1: <i>Ultomiris</i> Q8W</li> <li>• Arm 2: Placebo</li> </ul>	<ul style="list-style-type: none"> <li>• Primary endpoint: TMA response</li> <li>• Secondary endpoints: Time to TMA response, TMA relapse</li> </ul>	<ul style="list-style-type: none"> <li>• FPCD: Q4 2020</li> <li>• Data anticipated: &gt;2023</li> </ul>
Phase III ALXN-1210-TM-314 NCT04557735	Paediatric thrombotic microangiopathy associated with haematopoietic stem cell transplant	40	<ul style="list-style-type: none"> <li>• Arm 1: <i>Ultomiris</i> administered once every 4-8 weeks</li> </ul>	<ul style="list-style-type: none"> <li>• Primary endpoint: Proportion of participants with TMA response</li> <li>• Secondary endpoints: Time to TMA response, Proportion of participants with TMA relapse</li> </ul>	<ul style="list-style-type: none"> <li>• FPCD: Q4 2020</li> <li>• Data anticipated: &gt;2023</li> </ul>
Phase II ALXN-1210-NEPH-202 NCT04564339	Proliferative lupus nephritis or immunoglobulin A nephropathy	120	<ul style="list-style-type: none"> <li>• Arm 1: LN Cohort, <i>Ultomiris</i></li> <li>• Arm 2: LN Cohort, Placebo</li> <li>• Arm 3: IgAN Cohort, <i>Ultomiris</i></li> <li>• Arm 4: IgAN Cohort, Placebo</li> </ul>	<ul style="list-style-type: none"> <li>• Primary endpoint: Both cohorts, percentage change in proteinuria from baseline to Week 26</li> <li>• Secondary endpoints: Both cohorts, percentage change in proteinuria from baseline to Week 50</li> </ul>	<ul style="list-style-type: none"> <li>• FPCD: Q1 2021</li> <li>• Data anticipated: &gt;2023</li> </ul>



# Ultomiris (anti-complement C5 mAb)

## Neurology

Approved medicines

Late-stage development

Early development

Trial	Population	Patients	Design	Endpoints	Status
Phase III ALXN-1210-NMO-307 NCT04201262	Neuromyelitis optica spectrum disorder	58	<ul style="list-style-type: none"> <li>Arm 1: <i>Ultomiris</i> Q8W</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: time to first adjudicated on-trial relapse</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2019</li> <li>LPCD: Q1 2021</li> <li>Data anticipated: H1 2022</li> </ul>
Phase III ALXN-1210-MG-306 NCT03920293	Generalised myasthenia gravis	175	<ul style="list-style-type: none"> <li>Arm 1: <i>Ultomiris</i></li> <li>Arm 2: placebo</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: change from baseline in MG-ADL total score at Week 26</li> </ul>	<ul style="list-style-type: none"> <li>Data readout: Q2 2021</li> <li>Primary endpoint met</li> </ul>
Phase II/III ALXN-1210-ALS-308 NCT04248465	Amyotrophic lateral sclerosis	382	<ul style="list-style-type: none"> <li>Arm 1: <i>Ultomiris</i></li> <li>Arm 2: placebo</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: change from baseline in ALSFRS-R total score</li> </ul>	<ul style="list-style-type: none"> <li>Trial halted post futility analysis Q3 2021</li> </ul>
Phase II/III ALXN1210-DM-310 NCT04999020	Dermatomyositis	180	<ul style="list-style-type: none"> <li>Arm 1: <i>Ultomiris</i></li> <li>Arm 2: placebo</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint : improvement response on International Myositis Assessment And Clinical Studies-Total Improvement Score (IMACS-TIS)</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2021</li> <li>Data anticipated: &gt;2023</li> </ul>

Oncology

CVRM

R&I

Other

V&I

Rare Disease



# *Koselugo* (selumetinib, MEK inhibitor)

## Neurofibromatosis Type 1 (NF1)

Trial	Population	Patients	Design	Endpoints	Status
Phase III KOMET NCT04924608	Adult age ≥18 years with NF1 who have symptomatic, inoperable PN. Available baseline chronic target PN pain score	146	<ul style="list-style-type: none"> <li>Multicentre, international study with a parallel, randomized, double blind, placebo controlled, 2 arm design</li> <li>Arm A: <i>Koselugo</i> 25mg/m<sup>2</sup> BID</li> <li>Arm B: Placebo BID until end of Cycle 12, then cross over to <i>Koselugo</i> 25mg/m<sup>2</sup> BID</li> </ul>	<ul style="list-style-type: none"> <li>Primary Endpoint: ORR by end of Cycle 16 on <i>Koselugo</i> vs placebo as determined by ICR per REiNS criteria.</li> <li>Key secondary endpoint: Chronic PN-pain intensity (change from baseline) on <i>Koselugo</i> vs placebo</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2021</li> <li>Data anticipated: &gt;2023</li> </ul>



# Koselugo (selumetinib, MEK inhibitor)

## Paediatric neurofibromatosis type 1, solid tumours

Trial	Population	Patients	Design	Endpoints	Status
Phase II SPRINT NCT01362803 Partnered	Paediatric NF1	75	<ul style="list-style-type: none"> <li>Single arm: <i>Koselugo</i> 25mg/m<sup>2</sup> BID with 2 strata:</li> <li>Stratum 1: PN related morbidity present at enrolment</li> <li>Stratum 2: no PN related morbidity present at enrolment</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: Complete partial and complete response rate measured by volumetric MRI, Duration of response and functional outcomes/QoL</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2015</li> <li>LPCD: Q4 2016</li> <li>Data readout: Q1 2019</li> <li>Primary endpoint met</li> </ul>
Phase I/II SPRINKLE NCT05309668	Child age 1-6 years diagnosed with NF1 with symptomatic, inoperable PN Must have at least one measurable PN, defined as a PN of at least 3 cm measured in one dimension	38	<ul style="list-style-type: none"> <li>Single-Arm - <i>Koselugo</i></li> <li>Open Label</li> </ul>	<ul style="list-style-type: none"> <li>Primary Endpoints</li> <li>- <i>Koselugo</i> AUC0-12 derived after single dose administration [ Time Frame: Pre-dose and 1, 2, 3, 4, 6, 8 and 10-12 hours after <i>Koselugo</i> single dose on the first day of study treatment (Cycle 1 Day 1) (each cycle is 28 days) ]</li> <li>- Adverse Events graded by CTCAE Ver 5.0 [ Time Frame: from screening until 30 days after last dose ]</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2022</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase I Japan PK / Safety trial NCT04495127 Partnered	Paediatric inoperable NF1-PN patients	44904	<ul style="list-style-type: none"> <li>Open-label trial</li> <li><i>Koselugo</i> in Japanese paediatric NF1-PN patients</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: safety</li> <li>Secondary endpoints: PK, anti-tumour effect</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2020</li> <li>LPCD: Q4 2020</li> <li>Data readout: Q4 2021</li> </ul>
Phase I China PK / Safety / Efficacy trial NCT04590235	Paediatric (2-17 years old), adult NF1	32	<ul style="list-style-type: none"> <li>Single arm trial with 3 phases;</li> <li>Dose confirmation phase (n=6 for 3 cycles),</li> <li>Expansion phase (24mths post LSD)</li> <li>Long term follow up (60mths post LSD)</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: safety, tolerability and PK</li> <li>Secondary endpoints: efficacy (ORR, DoR; TTR; PFS)</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2020</li> </ul>
Phase I NCT05101148	Adolescents aged ≥ 12 to < 18 years at trial entry with a clinical diagnosis of NF1 related PN. <i>Koselugo</i> with a low-fat meal compared to fasted state	20	<ul style="list-style-type: none"> <li>Single-arm, multiple dose, sequential, two or three period trial</li> <li><i>Koselugo</i> 25mg/m<sup>2</sup> BID given with a low-fat meal versus the same dose given in a fasted state.</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoints: PK (steady state systemic exposure), safety (especially GI toxicity)</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2021</li> </ul>



# Acoramidis (ALXN2060)

## ATTR-CM

Approved medicines

Late-stage development

Early development

Trial	Population	Patients	Design	Endpoints	Status
Phase III ALXN2060-TAC-302 NCT04622046	ATTR-CM	22	<ul style="list-style-type: none"><li>Arm 1: 800 mg acoramidis (ALXN2060) administered twice daily</li></ul>	<ul style="list-style-type: none"><li>Primary endpoint: Change from Baseline to Month 12 of Treatment In Distance Walked During the Six-minute Walk Test, All-cause mortality and cardiovascular-related hospitalization over a 30-month period</li></ul>	<ul style="list-style-type: none"><li>FPCD: Q4 2020</li><li>Data anticipated: &gt;2023</li></ul>

Oncology

CVRM

R&I

Other

V&I

Rare Disease



# ALXN1840 (bis-choline tetrathiomolybdate)

## Wilson disease

Trial	Population	Patients	Design	Endpoints	Status
Phase III WTX101-301 FoCus NCT03403205	Wilson Disease	215	<ul style="list-style-type: none"> <li>Arm 1: ALXN1840 was administered orally for 48 weeks at doses ranging from 15 milligrams (mg) every other day (QOD) up to a titrated dose of 60 mg daily.</li> <li>Arm 2: Standard of care</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: daily mean AUEC of dNCC</li> <li>Secondary endpoint: change from baseline in the UWDRS Part II total score</li> </ul>	<ul style="list-style-type: none"> <li>Data readout: Q3 2021</li> <li>Primary endpoint met</li> </ul>
Phase II ALXN1840-WD-204 NCT04573309	Wilson Disease	10	<ul style="list-style-type: none"> <li>Arm 1: Participants will be administered ALXN1840 at a dose of 15 milligrams (mg)/day on Day 1 through Day 28 and then increased to 30 mg/day on Day 29 through Day 39</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: mean daily copper balance</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2020</li> <li>Data anticipated: H2 2022</li> </ul>
Phase II ALXN1840-WD-205 NCT04422431	Wilson Disease	31	<ul style="list-style-type: none"> <li>Arm 1: ALXN1840, Participants will be initiated at 15 milligrams once daily, then the dose will be increased to 30 milligrams once daily at Week 6.</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: change from baseline at Week 48 in liver copper concentration</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2020</li> <li>Data anticipated: H2 2022</li> </ul>



# CAEL-101 (fibril-reactive mAb)

## AL amyloidosis

Trial	Population	Patients	Design	Endpoints	Status
Phase III CAEL101-302 NCT04512235	Mayo stage IIIa amyloidosis	267	<ul style="list-style-type: none"> <li>Arm 1: CAEL-101 combined with SoC for plasma cell dyscrasia (PCD)</li> <li>Arm 2: placebo combined with SoC for PCD</li> </ul>	<ul style="list-style-type: none"> <li>Primary: time from first dose of trial drug until death or end of trial</li> <li>Secondary: change in distance walked during a six-minute walk test, and quality of life measures</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q4 2020</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase III CAEL101-301 NCT04504825	Mayo stage IIIb amyloidosis	110	<ul style="list-style-type: none"> <li>Arm 1: CAEL-101 combined with SoC for PCD</li> <li>Arm 2: placebo combined with SoC for PCD</li> </ul>	<ul style="list-style-type: none"> <li>Primary: time from first dose of trial drug until death or end of trial</li> <li>Secondary: change in distance walked during a six-minute walk test, and quality of life measures</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2021</li> <li>Data anticipated: &gt;2023</li> </ul>
Phase II CAEL101-203 NCT04304144	Mayo Stage I, Stage II and Stage IIIa amyloidosis	25	<ul style="list-style-type: none"> <li>Arm 1: CAEL-101 combined with SoC CyBorD</li> <li>Arm 2: placebo combined with SoC CyBorD and daratumumab</li> </ul>	<ul style="list-style-type: none"> <li>Primary: dose toxicity – occurrence of dose limiting toxicity (DLT) during the first 4 weeks of therapy</li> <li>Secondary: area under the plasma curve concentration versus time curve (AUC)</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2020</li> <li>Data anticipated: 2023</li> </ul>



# Danicopan (ALXN2040 factor D inhibitor)

## Haematology & ophthalmology

Trial	Population	Patients	Design	Endpoints	Status
Phase III ALXN2040-PNH-301 NCT04469465	PNH with clinically relevant extravascular haemolysis	84	<ul style="list-style-type: none"> <li>Arm 1: danicopan (ALXN2040) + C5 Inhibitor</li> <li>Arm 2: placebo + C5 Inhibitor</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: change from baseline in haemoglobin at week 12</li> <li>Secondary endpoint: percentage of participants with transfusion avoidance</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q1 2021</li> <li>Data anticipated: 2023</li> </ul>
Phase II ALXN2040-GA-201 NCT05019521	Geographic atrophy	330	<ul style="list-style-type: none"> <li>Arms 1-3: danicopan (ALXN2040) dosed at 100-400mg daily</li> <li>Arm 4: placebo</li> </ul>	<ul style="list-style-type: none"> <li>Primary endpoint: mean rate of change from baseline at week 52 in the square root of total GA lesion area in the trial eye as measured by FAF</li> </ul>	<ul style="list-style-type: none"> <li>FPCD: Q3 2021</li> <li>Data anticipated: &gt;2023</li> </ul>



Rare Disease –  
early-stage development



# ALXN1720 (anti-C5 bispecific minibody)

## Neurology

Approved medicines

Late-stage development

Early development

Trial	Population	Patients	Design	Endpoints	Status
Phase I ALXN1720-HV-101 NCT04920370	Healthy Volunteers	96	<ul style="list-style-type: none"><li>• Single &amp; Multiple Ascending Dosing of ALXN1720 via s.c. and i.v. administration</li><li>• Placebo</li></ul>	<ul style="list-style-type: none"><li>• Primary: Incidence of TEAEs and SAEs</li></ul>	<ul style="list-style-type: none"><li>• FPCD: Q3 2019</li><li>• Data anticipated: H1 2022</li></ul>

Oncology

CVRM

R&I

Other

V&I

Rare Disease



# ALXN1820 (anti-properdin)

## Haematology

Approved medicines  
Late-stage development  
Early development

Trial	Population	Patients	Design	Endpoints	Status
Phase I ALXN1820-HV-101 NCT04631562	Healthy Volunteers	80	<ul style="list-style-type: none"><li>• Arm 1: ALXN1820 administered s.c. or i.v., multiple ascending doses</li><li>• Arm 2: Placebo</li></ul>	<ul style="list-style-type: none"><li>• Primary: Participants with TEAEs</li></ul>	<ul style="list-style-type: none"><li>• FPCD: Q1 2021</li><li>• Data anticipated: H2 2022</li></ul>

Oncology

CVRM

R&I

Other

V&I

Rare Disease



# ALXN1850 (next generation asfotase alfa)

## Hypophosphatasia

Approved medicines

Late-stage development

Early development

Trial	Population	Patients	Design	Endpoints	Status
Phase I ALXN1850-HPP-101 NCT04980248	HPP	15	<ul style="list-style-type: none"><li>Arm 1: ALXN1850, 3 cohorts at low, medium, and high dosages</li></ul>	<ul style="list-style-type: none"><li>Primary: Incidence of TEAEs and TESAEs</li></ul>	<ul style="list-style-type: none"><li>FPCD: Q3 2021</li><li>Data anticipated: H2 2022</li></ul>

Oncology

CVRM

R&I

Other

V&I

Rare Disease



# ALXN2050 (factor D inhibitor)

Haematology, nephrology, neurology

Approved medicines

Late-stage development

Early development

Trial	Population	Patients	Design	Endpoints	Status
Phase II ACH228-110 NCT04170023	PNH	26	<ul style="list-style-type: none"><li>Arm 1: ALXN2050 Monotherapy, with groups including treatment-naïve, C5 inhibitor treatment experienced, and patients previously receiving danicopan (ALXN2040)</li></ul>	<ul style="list-style-type: none"><li>Primary endpoint: Change in haemoglobin relative to baseline</li><li>Secondary endpoints: Number of participants who have transfusion avoidance, Change in lactate dehydrogenase relative to baseline</li></ul>	<ul style="list-style-type: none"><li>FPCD: Q4 2019</li><li>Data anticipated: &gt;2023</li></ul>

Oncology

CVRM

R&I

Other

V&I

Rare Disease



# List of abbreviations

14C	Radioactive isotope of carbon, Carbon 14
1L, 2L, 3L	1st, 2nd or 3rd line
5-FU	5-fluorouracil
A2AR	Adenosine A2A receptor
ACQ	Asthma control questionnaire
ACR	American college of rheumatology response scoring system
ADA	Anti-drug antibodies
ADC	Antibody-drug conjugate
ADP	Adenosine diphosphate
AE	Adverse event
aHUS	Atypical haemolytic uraemic syndrome
AI	Auto-injector
AKT	Protein kinase B
ALK	Anaplastic large-cell lymphoma kinase
ALSFRS-R	Amyotrophic Lateral Sclerosis Functional Rating Scale-Revised
APFS	Accessorised pre-filled syringe
AQLQ	Asthma quality of life questionnaire
AS	Albuterol sulphate
ATM	Ataxia-telangiectasia mutated kinase
ATR	Ataxia telangiectasia and rad3-related protein
ATTR-CM	Transthyretin amyloid cardiomyopathy
AUC	Area under curve
AUEC	Area under the effect-time curve
B7RP	B7-related protein-1
BA	Bioavailability
BAFF	B-cell activating factor
BCG	Bacillus Calmette–Guérin
BCMA	B-cell maturation antigen
BDA	Budesonide albuterol
BFF	Budesonide and formoterol fumarate
BGF	Budesonide, glycopyrronium and formoterol fumarate
BICR	Blinded independent central review
BID	Bis in die (twice per day)
BIG	Big ten cancer research consortium
BMD	Bone mineral density
BMFI	Bone metastasis-free interval
BMI	Body mass index
BRCAwt	Breast cancer wild-type gene
BRD4	Bromodomain-containing protein 4
BTC	Biliary tract carcinoma

BTK	Bruton's tyrosine kinase
CA-125	Cancer antigen 125
CAD	Coronary artery disease
CBR	Clinical benefit rate
CD	Cluster of differentiation
CDK	Cyclin-dependent kinase
CE	Clinically evaluable
CHD	Coronary heart disease
Chemo	Chemotherapy
CHF	Chronic heart failure
CKD	Chronic kidney disease
CLL	Chronic lymphocytic leukaemia
CMAX	Maximum observed plasma concentration
C-MET	Tyrosine-protein kinase Met
CNS	Central nervous system
COPD	Chronic obstructive pulmonary disease
CR	Complete response
CRC	Colorectal cancer
CrCl	Creatinine clearance
CRR	Complete response rate
CTC	Circulating tumour cell
CTLA-4	Cytotoxic T-lymphocyte-associated antigen 4
CV	Cardiovascular
CVOT	Cardiovascular outcomes trial
CVRM	Cardiovascular renal and metabolism
CXCR2	C-X-C Motif chemokine receptor 2
DB	Double blind
DC	Disease control
DCR	Disease control rate
DDI	Drug-drug Interaction
dECG	Differentiated electrocardiogram
DFS	Disease free survival
DLBCL	Diffuse large B-cell lymphoma
DLT	Dose-limiting toxicity
DMARDs	Disease-modifying antirheumatic drugs
DNA	Deoxyribonucleic acid
DoCR	Durability of complete response
dNCC	Directly Measured Non-ceruloplasmin-bound Copper
DoR	Duration of response
DPI	Dry powder inhaler

DRFI	Disease recurrence-free interval
DXA	Dual energy X-ray absorptiometry
EBRT	External beam radiation therapy
ECG	Electrocardiogram
EFS	Event-free survival
eGFR	Estimated glomerular filtration rate
EGFR	Epidermal growth factor receptor
ER	Oestrogen receptor
ERK	Extracellular signal-regulated kinase
ESCC	Esophageal squamous cell carcinoma
ESR	Externally sponsored trial
ESR1	Oestrogen receptor 1
ET	Endocrine therapy
FAF	Fundus Autofluorescence
FDC	Fixed-dose combination
FeNO	Fractional nitric oxide concentration in exhaled breath
FEV	Forced-expiratory volume
FGFR	Fibroblast growth factor receptor
FLAP	5-lipoxygenase-activating protein
FPDC	First patient commenced dosing
FPG	Fasting plasma glucose
GA	Gestational age
GA	Geographic atrophy
GBM	Glioblastoma
gBRCAm or tBRCAm	Germline or tumour (somatic) BRCA mutation
GEJ	Gastric/gastro-oesophageal junction
GFF	Glycopyrronium and formoterol fumarate
GLP-1	Glucagon-like peptide-1
GMFRs	Geometric mean fold rises
GMTs	Geometric mean titers
hADME	Human mass balance
HAI	Haemagglutination-inhibition
HbA1c	Haemoglobin A1c
HCC	Hepatocellular carcinoma
HD	High dose
HDL-C	High-density lipoprotein cholesterol
HER2	Human epidermal growth factor receptor 2
HF	Heart failure
HFpEF	Heart failure with preserved ejection fraction



# List of abbreviations

HFrEF	Heart failure with reduced ejection fraction
HGFR	Met/hepatocyte growth factor receptor
HGSC	High grade serous carcinoma
hHF	Hospitalisation for heart failure
HIF-PHI	Hypoxia inducible factor - prolyl hydroxylase inhibitor
HNSCC	Head and neck squamous-cell carcinoma
HPV	Human papillomavirus
HRD	Homologous recombination deficiency
HRRm	Homologous recombination repair mutation
i	inhibitor
IA	Investigator-assessed
ICS	Inhaled corticosteroid
ICU	Intensive care unit
IDFS	Invasive disease-free survival
IgAN	Immunoglobulin A nephropathy
IL	Interleukin
i.m.	Intramuscular
IRC	Independent review committee
ISS	Investigator-sponsored studies
i.v.	Intravenous
J-SD	Japanese single dose
Ki67	Protein that is encoded by the MKI67 gene in human
LAAB	Long-acting antibody
LABA	Long -acting beta agonist
LAMA	Long-acting muscarinic agonist
LCAT	Lecithin-cholesterol acyltransferase
LCM	Lifecycle management
LDH	Lactate dehydrogenase
LN	Lupus nephritis
LOCS III	Lens opacities classification system III
LPCD	Last patient commenced dosing
LV	Left ventricle
m	Mutation
mAb	Monoclonal antibody
MABA	Muscarinic antagonist-beta2 agonist
MACE	Major adverse cardiac events
MAD	Multiple ascending dose
MCC	Mucociliary clearance
MCL	Mantle cell lymphoma
MCL1	Myeloid leukemia cell differentiation protein 1
mCRPC	Metastatic castrate-resistant prostate cancer
MD	Medium dose
MDI	Metered-dose inhaler
MDS	Myelodysplastic syndrome
MEK	Mitogen-activated protein kinase
MET	Tyrosine-protein kinase Met
MG-ADL	Myasthenia Gravis-Activities Of Daily Living
MI	Myocardial infarction
MMT	Mixed meal test
MPO	Myeloperoxidase
mPR	Major pathological response
MRI	Magnetic resonance imaging
MTD	Maximum tolerated dose
NaC	Sodium channel
NCI	National cancer institute (US)
NCPV	Noncalcified plaque volume
NF1	Neurofibromatosis type 1
NGF	Nerve growth factor
NHL	Non-Hodgkin's lymphoma
NIH	National Institute of Health (US)
NKG2a	Natural killer cell C-type lectin receptor G2A
NME	New molecular entity
NRG	National clinical trials network in oncology (US)
NSCLC	Non-small cell lung cancer
OCS	Oral corticosteroid
OD	Once daily
OGTT	Oral glucose tolerance test
OR	Objective response
ORR	Objective response rate
OS	Overall survival
PARP	Poly ADP ribose polymerase
PASI	Psoriasis area severity index
PBD	Pyrrrolbenzodiazepine
pCR	Pathological complete response
PD	Pharmacodynamics
PD-1	Programmed cell death protein 1
PDAC	Pancreatic ductal adenocarcinoma
PDE4	Phosphodiesterase type 4
PD-L1	Programmed death-ligand 1
PET	Positron-emission tomography
PFS	Progression free survival
PgR	Progesterone receptor
PI3K	Phosphoinositide 3-kinase
PIK3CA	Phosphatidylinositol 3 kinase catalytic alpha gene
PK	Pharmacokinetics
PLL	Prolymphocytic leukaemia
pMDI	Pressurised metered dose inhaler
PN	Plexiform neurofibromas
PNH	Paroxysmal nocturnal haemoglobinuria
POC	Proof of concept
POM	Proof of mechanism
pPCI	Primary percutaneous coronary intervention
PR	Partial response
pre-BD	Pre-bronchodilator
PRO	Patient reported outcome
PRR	Recurrent platinum resistant
PS	Propensity score
PSA	Prostate-specific antigen
PSC	Pulmonary sarcomatoid carcinoma
PSMA	Prostate-specific membrane antigen
PTEN	Phosphatase and tensin homolog gene
Q2,3,4,8W	Quaque (every) two, three... weeks
QD	Quaque in die (once a day)
QID	Quarter in die (four times a day)
QOD	Quaque altera die (every other day)
QoL	Quality of Life
QTcF	Corrected QT interval by Fredericia
RA	Rheumatoid Arthritis
RAAS	Renin-angiotensin-aldosterone system
RECIST	Response evaluation criteria in solid tumours
RFS	Relapse-free survival
rhLCAT	Recombinant human Lecithin-cholesterol acyltransferase
RORγ	Related orphan receptor gamma
r/r	Relapsed/refractory
RSV	Respiratory Syncytial Virus
RT	Radiation therapy
R&I	Respiratory and Immunology
SABA	Short-acting beta2-agonist
SAD	Single ascending dose
SAE	Serious adverse event



# List of abbreviations

SBRT	Stereotactic body radiation therapy
s.c.	Subcutaneous
SCCHN	Squamous-cell carcinoma of the head and neck
SCLC	Small cell lung cancer
SD	Stable disease
SERD	Selective oestrogen receptor degrader
SGLT2	Sodium-glucose transport protein 2
SGRM	Selective glucocorticoid receptor modulator
SGRQ	Saint George respiratory questionnaire
SJC	Swollen joint count
SLE	Systemic lupus erythematosus
SLL	Small lymphocytic lymphoma
SMAD	Single and multiple ascending dose trial
SoC	Standard of care
sPGA	Static physician's global assessment score
STAT3	Signal transducer and activator of transcription 3
sUA	Serum uric acid
T2DM	Type 2 Diabetes Mellitus
T790M	Threonine 790 substitution with methionine
TACE	Transarterial Chemoembolization
TEAEs	Treatment-emergent adverse events
TESAEs	Treatment-emergent serious adverse events
TID	Ter in die (three times a day)
TJC	Tender joint count
TKI	Tyrosine kinase inhibitor
TLR	Toll-like receptor 9
TMA	Thrombotic microangiopathy
TNBC	Triple negative breast cancer
TNF	Tumour necrosis factor
TSLP	Thymic stromal lymphopietin
TTF	Time to treatment failure
TTNT	Time to next therapy
TTP	Time to tumour progression
UACR	Urine albumin creatinine ratio
UMEC	Umeclidinium
URAT1	Uric Acid Transporter 1
UWDRS	Unified Wilson Disease Rating Scale
VEGF	Vascular endothelial growth factor
V&I	Vaccine & Immune therapies
YTE	Triple-amino-acid (M252Y/S254T/T256E [YTE]) substitution

